

Modules and Categories*

Lenny Taelman



*Preliminary version, February 27, 2018

Contents

Foreword	7
Prerequisites	7
Other sources	7
Acknowledgements	7
Chapter 1. Modules over a ring	9
1. Left and right modules	9
2. First examples	10
3. Homomorphisms, submodules and quotient modules	12
4. Products, direct sums and free modules	13
Exercises	17
Chapter 2. Exact sequences	21
1. Exact sequences	21
2. The Five Lemma and the Snake Lemma	22
3. Split short exact sequences	24
Exercises	26
Chapter 3. Finitely generated modules over a PID	31
1. Introduction	31
2. Review of principal ideal domains	31
3. Free modules of finite rank over a PID	32
4. Structure of finitely generated modules over a PID	33
5. Application to Jordan normal form	35
Exercises	38
Chapter 4. Categories	41
1. Definition	41
2. Big examples	42
3. Small examples	42
4. Isomorphisms, monomorphisms, epimorphisms	44

5. Initial and final objects	45
Exercises	47
Chapter 5. Functors	51
1. Definition of a functor	51
2. Many examples	51
3. Contravariant functors	54
4. Functors with multiple arguments	55
5. Morphisms of functors	56
6. Equivalences of categories	57
Exercises	61
Chapter 6. Tensor product	65
1. Tensor product of a right and a left module	65
2. Tensor products and bimodules	68
3. Tensor product as a functor	70
4. The adjunction	72
5. Restriction and extension of scalars	73
Exercises	75
Chapter 7. Adjoint functors	79
1. Adjoint pairs of functors	79
2. Many examples	80
3. Yoneda and uniqueness of adjoints	83
Exercises	86
Chapter 8. Limits and colimits	89
1. Product and coproduct	89
2. Pushout and pullback	92
3. Equalizer and co-equalizer	95
4. Limits and colimits	96
5. Yoneda and limits and colimits of sets	99
6. Adjoint functors and limits	101
Exercises	103
Chapter 9. Chain complexes	107
1. Chain complexes and their homology modules	107
2. The long exact sequence	108
3. The homotopy category	109
Exercises	112

Chapter 10. Free resolutions	115
1. Definition and existence	115
2. The free resolution functor	117
Exercises	121
Chapter 11. The Ext functors	123
1. The functors Ext^n	123
2. The long exact sequence	125
3. Ext^1 and extensions	126
Exercises	129
Bibliography	133

Foreword

These are course notes for a one-semester third-year course on categories and modules taught at the University of Amsterdam.

Prerequisites

Linear algebra, groups, elementary point-set topology, and the basics of rings and fields. Several exercises and examples in the parts about categories refer to basic concepts in algebraic topology, Galois theory, or representation theory. Although in principle these can be skipped, developing a rich vocabulary of natural examples is probably the most important aspect of becoming acquainted with categories.

Other sources

Excellent alternative sources for much of the material in this course are the concise but clear and well-written Atiyah & MacDonald [1], the more lengthy Lang [2], the course notes by Moerdijk [3] and (in Dutch) the Algebra 2 course notes by Stevenhagen [4]. A good introduction in the language of categories and functors containing much more than these notes is [5].

Acknowledgements

Many thanks to Wessel Bindt, David de Boer, Jeroen Dekker, Koen van Duin, Eline Filius, Pepijn Hoefgeest, Tycho van Hoof, Juultje Kok, Christopher Spelt, Tijn de Vos, for their many corrections and suggestions to the first drafts of these course notes.

CHAPTER 1

Modules over a ring

1. Left and right modules

Let R be a ring. Recall that this means R is a set equipped with an addition $(s, t) \mapsto s + t$, multiplication $(s, t) \mapsto st$, and distinguished elements $0 \in R$ and $1 \in R$ satisfying

- (R1) $(R, +, 0)$ is an abelian group,
- (R2) for all $r, s, t \in R$ we have $(rs)t = r(st)$,
- (R3) for all $r, s, t \in R$ we have $r(s+t) = rs+rt$ and $(r+s)t = rt+st$,
- (R4) for all $r \in R$ we have $1r = r1 = r$.

DEFINITION 1.1. A *left module* over a ring R is an abelian group M equipped with an operation

$$R \times M \rightarrow M, (r, x) \mapsto rx$$

satisfying for all $r, s \in R$ and $x, y \in M$ the following identities:

- (M1) $r(x + y) = rx + ry$,
- (M2) $(r + s)x = rx + sx$,
- (M3) $(rs)x = r(sx)$,
- (M4) $1x = x$.

One also says that ‘ R acts on M ’, so that axiom (M3) for example expresses that acting by rs is the same as first acting by s , and then by r .

We use the same symbol 0 to denote the elements $0 \in M$ and $0 \in R$. This should not lead to confusion, see Exercise 1.1.

A *right module* over R is defined similarly: the action is written on the right: $M \times R \rightarrow M, (x, r) \mapsto xr$, and must satisfy

- (M1') $(x + y)r = xr + yr$,
- (M2') $x(r + s) = xr + xs$,
- (M3') $x(rs) = (xr)s$,
- (M4') $x1 = x$.

If R is commutative, then the difference between a left and a right module is purely a matter of notation, but over a non-commutative ring the axioms (M3) and (M3') give genuinely different conditions.

We will mostly work with left modules, and simply call them *modules over R* or *R -modules*.

There is another way to describe (left) modules, using the endomorphism ring of an abelian group. Let A be an abelian group. Denote by $\text{End}(A)$ the set of group homomorphisms $A \rightarrow A$. This forms a ring with addition and multiplication of $f, g \in \text{End}(A)$ defined by pointwise addition

$$f + g: A \rightarrow A, a \mapsto f(a) + g(a)$$

and composition

$$fg: A \rightarrow A, a \mapsto f(g(a)).$$

The zero element of this ring is the constant map $0: a \mapsto 0$, and the unit element is the identity map $\text{id}_A: a \mapsto a$.

LEMMA 1.2. *Let M be an R -module. Then the map*

$$R \rightarrow \text{End}(M), r \mapsto (x \mapsto rx)$$

is a ring homomorphism. Conversely, let M be an abelian group and $\phi: R \rightarrow \text{End}(M)$ be a ring homomorphism. Then the operation

$$R \times M \rightarrow M, (r, x) \mapsto rx := \phi(r)(x)$$

gives M the structure of an R -module.

PROOF. See Exercise 1.3. □

In other words, a (left) R -module is the same as an abelian group M together with a ring homomorphism $R \rightarrow \text{End}(M)$.

2. First examples

EXAMPLE 1.3. Let R be a ring. Then the trivial group $\{0\}$ is an R -module with $r0 := 0$ for all $r \in R$. We denote this module by 0 , and call it the *zero module*.

EXAMPLE 1.4. Let R be a ring and $n \geq 0$. Then $M := R^n$ is an R -module with addition

$$(x_1, \dots, x_n) + (y_1, \dots, y_n) := (x_1 + y_1, \dots, x_n + y_n)$$

and R -action

$$r \cdot (x_1, \dots, x_n) := (rx_1, \dots, rx_n).$$

For $n = 0$ we obtain the zero module $R^0 = 0$ and for $n = 1$ we obtain the R -module R .

EXAMPLE 1.5. For every abelian group A there is a unique ring homomorphism $\mathbf{Z} \rightarrow \text{End}(A)$. It follows that a \mathbf{Z} -module is the same as an abelian group. For $r \in \mathbf{Z}$ and $x \in A$ we have

$$r \cdot x = \begin{cases} x + \cdots + x & (r \text{ terms}) & r \geq 0 \\ -(x + \cdots + x) & (-r \text{ terms}) & r \leq 0 \end{cases}$$

EXAMPLE 1.6. Let K be a field. Then a K -module is the same as a K -vector space.

EXAMPLE 1.7. Let K be a field and $n \geq 0$. Let $\text{Mat}_n(K)$ be the ring of n by n matrices over K . Then K^n is a left $\text{Mat}_n(K)$ -module via

$$\text{Mat}_n(K) \times K^n \mapsto K^n, (A, v) \mapsto A \cdot v,$$

where we interpret vectors $v \in K^n$ as column matrices.

EXAMPLE 1.8. Let R be a ring and $I \subset R$ an ideal. Then I is an R -module.

EXAMPLE 1.9. Let K be a field. Let V be a K -vector space, and $\alpha: V \rightarrow V$ be a K -linear endomorphism of V . Then there is a unique ring homomorphism

$$\rho: K[X] \rightarrow \text{End}(V)$$

such that

- (1) $\rho(\lambda)(v) = \lambda v$ for all $\lambda \in K$ and $v \in V$,
- (2) $\rho(X) = \alpha$.

This homomorphism is given by

$$(1) \quad \rho: \sum \lambda_i X^i \mapsto \left(v \mapsto \sum \lambda_i \alpha^i(v) \right),$$

where α^i denotes the iterated composition $\alpha \circ \cdots \circ \alpha$. In particular, V obtains the structure of a $K[X]$ -module.

Conversely, given a $K[X]$ -module V , the restriction of the action of $K[X]$ to K makes V into a K -vector space, and the map

$$\alpha: V \rightarrow V, v \mapsto X \cdot v$$

is K -linear. We conclude that a $K[X]$ -module is the same as a K -vector space equipped with an endomorphism (namely the action of X).

We will see in Chapter 3 that the ‘Jordan normal form’ of complex square matrices (\mathbf{C} -linear endomorphisms of \mathbf{C}^n), is really a theorem about the structure of $\mathbf{C}[X]$ -modules, and that it is most naturally explained in terms of ideals in $\mathbf{C}[X]$.

EXAMPLE 1.10 (The group algebra and representations). Let K be a field and G a group. Let $K[G]$ be the group algebra of G over K . Elements of $K[G]$ are formal expressions

$$\sum_{g \in G} a_g g \quad (a_g \in K)$$

with $a_g = 0$ for all but finitely many g (this is automatic if G is a finite group). Addition is defined in the obvious way. Multiplication is defined by extending the multiplication in G . We have

$$\left(\sum_{g \in G} a_g g \right) \cdot \left(\sum_{h \in G} b_h h \right) = \sum_{t \in G} c_t t$$

with

$$c_t = \sum_{gh=t} a_g b_h.$$

A $K[G]$ -module is the same as a K -vector space V , together with a group homomorphism

$$G \rightarrow \mathrm{GL}_K(V) = \mathrm{End}_K(V)^\times.$$

In other words, a $K[G]$ -module is a K -linear representation of G .

3. Homomorphisms, submodules and quotient modules

DEFINITION 1.11. Let M and N be R -modules. An R -module homomorphism from M to N is a map $f: M \rightarrow N$ such that for all $r \in R$ and $x, y \in M$ we have

$$f(x + y) = f(x) + f(y)$$

and

$$f(rx) = rf(x).$$

We also say that the map $f: M \rightarrow N$ is R -linear. The set of R -module homomorphisms from M to N is denoted $\mathrm{Hom}_R(M, N)$. An *isomorphism* of R -modules is a bijective R -module homomorphism. Two R -modules are called *isomorphic* if there exists an isomorphism between them.

A *submodule* of an R -module M is a subgroup $N \subset M$ such that for all $r \in R$ and $x \in N$ we have $rx \in N$. A submodule of an R -module is itself an R -module. If $N \subset M$ is a submodule, then the abelian group M/N has the structure of an R -module, via

$$r(x + N) := rx + N.$$

We call M/N the *quotient module*.

To a module homomorphism $f: M \rightarrow N$ are associated three important modules. The *kernel*

$$\ker f := \{x \in M \mid f(x) = 0\} \subset M,$$

which is a submodule of M , the *image*

$$\operatorname{im} f := f(M) \subset N,$$

which is a submodule of N , and the *cokernel*

$$\operatorname{coker} f := N/(\operatorname{im} f),$$

which is a quotient module of N . A homomorphism f is injective if and only if $\ker f$ is trivial, and it is surjective if and only if $\operatorname{coker} f$ is trivial.

As with groups or vector spaces, we have the natural isomorphism

$$M/(\ker f) \xrightarrow{\sim} \operatorname{im} f, \quad \bar{x} \mapsto f(x).$$

4. Products, direct sums and free modules

Let R be a ring and let M and N be R -modules. The cartesian product $M \times N$ is naturally an R -module with $(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 + y_2)$ and $r(x_1, x_2) = (rx_1, rx_2)$. We call this R -module the *product* or *direct product* of the R -modules M and N . More generally, if $(M_i)_{i \in I}$ is a family of R -modules indexed by a set I , then the product $\prod_{i \in I} M_i$ is naturally an R -module with

$$(x_i)_{i \in I} + (y_i)_{i \in I} = (x_i + y_i)_{i \in I}, \quad r(x_i)_{i \in I} = (rx_i)_{i \in I},$$

for all $(x_i)_{i \in I}, (y_i)_{i \in I}$ in $\prod_{i \in I} M_i$ and $r \in R$. The empty product gives the zero module.

The *direct sum* of a collection $(M_i)_{i \in I}$ of R -modules indexed by a set I , denoted $\bigoplus_{i \in I} M_i$ is the R -submodule of $\prod_{i \in I} M_i$ defined as

$$\bigoplus_{i \in I} M_i := \left\{ (x_i)_{i \in I} \in \prod_{i \in I} M_i \mid \{i \in I : x_i \neq 0\} \text{ is finite} \right\}.$$

Note that this is indeed a submodule: if $(x_i)_{i \in I}$ and $(y_i)_{i \in I}$ have only finitely many non-zero terms, then so do $(x_i + y_i)_{i \in I}$ and $(rx_i)_{i \in I}$.

One sometimes phrases the condition on $(x_i)_{i \in I}$ as ‘ x_i is zero for all but finitely many i ’. Of course, if I is finite then the condition is vacuous and we have $\bigoplus_{i \in I} M_i = \prod_{i \in I} M_i$.

We have natural ‘inclusion’ maps

$$\iota_j: M_j \rightarrow \bigoplus_{i \in I} M_i, \quad x \mapsto \iota_j(x), \quad \iota_j(x)_i = \begin{cases} x & i = j \\ 0 & i \neq j \end{cases}$$

and ‘projection’ maps

$$\pi_j: \prod_{i \in I} M_i \rightarrow M_j, \quad (x_i)_{i \in I} \mapsto x_j.$$

If we have $M_i = M$ for all i , then we write $M^I := \prod_{i \in I} M$ and $M^{(I)} := \bigoplus_{i \in I} M$, so that we have

$$M^I = \{(x_i)_{i \in I} \mid x_i \in M\}$$

and

$$M^{(I)} = \{(x_i)_{i \in I} \mid x_i \in M, \text{ and } x_i = 0 \text{ for all but finitely many } i\}.$$

If I is a finite set of cardinality n , then we have $M^I = M^{(I)} \cong M^n$.

Let $(x_i)_{i \in I}$ be a family of elements of an R -module M . Then we call the intersection of all submodules $N \subset M$ that contain all x_i the *submodule generated by $(x_i)_{i \in I}$* . We denote it by $\langle x_i \rangle_{i \in I}$. It is the smallest submodule of M containing all the x_i . It consists of all finite R -linear combinations of the x_i .

We say that M is *finitely generated* if there exists a finite family $(x_i)_{i \in I}$ with $M = \langle x_i \rangle$.

EXAMPLE 1.12. Let R be a ring and I a set. Consider the module

$$R^{(I)} = \{(r_i)_{i \in I} \in R^I \mid r_i = 0 \text{ for all but finitely many } i \in I\}.$$

For an index $i \in I$ we denote by $e_i \in R^{(I)}$ the element $e_i := \iota_i(1)$. One may think of e_i as the ‘standard basis’ element

$$e_i = (\dots, 0, 0, 1, 0, 0, \dots)$$

with a 1 at position i . Clearly, every element of $R^{(I)}$ is a finite R -linear combination of e_i ’s, so the family $(e_i)_{i \in I}$ generates $R^{(I)}$. Note that if I is infinite, then the e_i do not generate the direct product R^I .

PROPOSITION 1.13. *Let R be a ring, M an R -module, and $(x_i)_{i \in I}$ a family of elements of M . Then there exists a unique R -linear map $\varphi: R^{(I)} \rightarrow M$ with $\varphi(e_i) = x_i$ for every $i \in I$.*

Moreover, φ is surjective if and only if M is generated by $(x_i)_{i \in I}$.

This generalises the basic fact from linear algebra that giving a linear map $\mathbf{R}^n \rightarrow V$ is the same as giving the images of the standard basis vectors.

PROOF OF PROPOSITION 1.13. The map φ is given by

$$\varphi: R^{(I)} \rightarrow M, (r_i)_{i \in I} \mapsto \sum_{i \in I} r_i x_i$$

(note that in the sum only finitely many terms are non-zero). This map is surjective if and only if every element of M can be written as a finite R -linear combination of elements x_i . \square

DEFINITION 1.14. Let M be an R -module and $(x_i)_{i \in I}$ a family of elements of M . Let $\varphi: R^{(I)} \rightarrow M$ be the unique R -linear map with $\varphi(e_i) = x_i$ for all $i \in I$. We say that $(x_i)_{i \in I}$ is a *basis* of M if φ is an isomorphism. We say that an R -module M is *free* if it has a basis. If it has a basis of cardinality n , then we say that M is *free of rank n* .

In contrast with the case of vector spaces (modules over a field), a finitely generated R -module need not have a basis. For example: for $m > 1$ the \mathbf{Z} -module $\mathbf{Z}/m\mathbf{Z}$ does not have a basis, and hence is not free.

PROPOSITION 1.15. *Let M be an R -module and $(x_i)_{i \in I}$ a family of elements of M . Then $(x_i)_{i \in I}$ is a basis of M if and only if for every $x \in M$ there is a unique family $(r_i)_{i \in I}$ of elements in R with*

- (1) $r_i = 0$ for all but finitely many i , and
- (2) $x = \sum_{i \in I} r_i x_i$.

Note that the condition in (1) guarantees that only finitely many terms in the sum in (2) are non-zero.

PROOF OF PROPOSITION 1.15. This is a direct translation of the definition: existence of (r_i) is equivalent with x being in the image of $\varphi: R^{(I)} \rightarrow M$, and uniqueness is equivalent with $\varphi: R^{(I)} \rightarrow M$ being injective. \square

EXAMPLE 1.16. Let U be an open subset of \mathbf{R}^n . Then the space of 1-forms $\Omega^1(U)$ on U forms a module over the ring $\mathcal{C}^\infty(U)$ of \mathcal{C}^∞ functions on U . This module is free of rank n , with basis dx_1, \dots, dx_n .

PROPOSITION 1.17. *Let R be a commutative ring with $0 \neq 1$. If R^n and R^m are isomorphic R -modules, then $n = m$.*

In other words: an R -module over a non-zero commutative ring which is free of finite rank has a well-defined rank. We already know this if R is a field (any two bases of a vector space have the same cardinality), and the proof of the proposition will be by reduction to the case of a field.

PROOF OF PROPOSITION 1.17. Let $M = R^m$, $N = R^n$ and let

$$\varphi: M \rightarrow N$$

be an isomorphism. Let $I \subset R$ be a maximal ideal (which exists in every non-zero commutative ring). Then φ induces an isomorphism

$$\bar{\varphi}: M/IM \rightarrow N/IN$$

of R/I -modules (see also Exercise 1.11). We have $M/IM = (R/I)^m$ and $N/IN = (R/I)^n$. But R/I is a field, so using the fact that a vector space has a well-defined dimension we find

$$m = \dim_{R/I} M/IM = \dim_{R/I} N/IN = n,$$

which is what we had to prove. □

Exercises

EXERCISE 1.1. Let M be an R -module. Show that for every $r \in R$ and $x \in M$ the following identities in M hold:

- (1) $r0 = 0$,
- (2) $0x = 0$,
- (3) $(-r)x = r(-x) = -(rx)$.

EXERCISE 1.2. Let $R = \{0\}$ be the zero ring. Show that every R -module is the zero module.

EXERCISE 1.3. Prove Lemma 1.2.

EXERCISE 1.4. Let $R = (R, 0, 1, +, \cdot)$ be a ring. Consider the opposite ring $R^{\text{opp}} = (R, 0, 1, +, \cdot^{\text{opp}})$ where multiplication is defined by

$$r \cdot^{\text{opp}} s := s \cdot r.$$

Show that a *right module* over R is the same as an abelian group M equipped with a ring homomorphism $R^{\text{opp}} \rightarrow \text{End}(M)$.

EXERCISE 1.5. Let M be an R -module, and let x_1, \dots, x_n be elements of M . Verify that the map

$$R^n \rightarrow M, (r_1, \dots, r_n) \mapsto r_1x_1 + \dots + r_nx_n$$

is a homomorphism of R -modules.

EXERCISE 1.6. Let R be a domain and $I \subset R$ a non-zero principal ideal. Show that I , as an R -module, is isomorphic to the R -module R .

EXERCISE 1.7. Let R be a ring, and let M and N be R -modules. Show that point-wise addition makes $\text{Hom}_R(M, N)$ into an abelian group. Show that if R is commutative, then $\text{Hom}_R(M, N)$ has a natural structure of R -module.

EXERCISE 1.8. Let R be a ring. Show $\text{Hom}_R(R, M) \cong M$ as abelian groups (and if R is commutative, as R -modules).

EXERCISE 1.9. Verify that the map (1) in Example 1.9 is indeed a ring homomorphism.

EXERCISE 1.10. Let K be a field and n a positive integer. Let V be a K -vector space, and let $\alpha_1, \alpha_2, \dots, \alpha_n$ be pairwise *commuting* linear endomorphisms of V . Show that there is a unique ring homomorphism

$$\rho: K[X_1, \dots, X_n] \rightarrow \text{End}(V)$$

such that for all $\lambda \in K$ and $v \in V$ we have $\rho(\lambda)(v) = \lambda v$ and for all i we have $\rho(X_i)(v) = \alpha_i(v)$.

Convince yourself that a $K[X_1, \dots, X_n]$ -module is the same thing as a vector space together with n pairwise commuting linear endomorphisms.

EXERCISE 1.11. Let R be a ring and $I \subset R$ a (two-sided) ideal. Let M be an R -module. Show that

$$IM := \left\{ \sum r_i x_i \mid r_i \in I, x_i \in M \right\}$$

is a sub- R -module of M , and show that M/IM is an R/I -module. Show that if M is free of rank n as R -module, then M/IM is free of rank n as R/I -module.

EXERCISE 1.12. Let R be a ring and let M be a left R -module. Show that

$$\text{Ann}_R(M) := \{ r \in R \mid rx = 0 \text{ for all } x \in M \}$$

is a (two-sided) ideal in R .

EXERCISE 1.13. Show that \mathbf{Q} is not a finitely generated \mathbf{Z} -module.

EXERCISE 1.14. Let K be a field and I a countably infinite set. Show that the K -module K^I does not have a countable generating set.

EXERCISE 1.15. Show that \mathbf{Q} is not a free \mathbf{Z} -module.

EXERCISE 1.16. Prove the following generalisation of Proposition 1.13: Let R be a ring and let $(M_i)_{i \in I}$ be a collection of R -modules indexed by a set I . Let N be an R -module, and let $(f_i: M_i \rightarrow N)_i$ be a collection of R -linear maps. Then there exists a unique R -linear map

$$f: \bigoplus_{i \in I} M_i \rightarrow N$$

such that for all $i \in I$ and $x \in M_i$ we have $f(\iota_i(x)) = f_i(x)$.

EXERCISE 1.17. Let R be a ring, $(M_i)_{i \in I}$ a family of R -modules, and N an R -module. Show that there are isomorphisms

$$\text{Hom}_R\left(\bigoplus_{i \in I} M_i, N\right) \xrightarrow{\sim} \prod_{i \in I} \text{Hom}_R(M_i, N)$$

and

$$\text{Hom}_R\left(N, \prod_{i \in I} M_i\right) \xrightarrow{\sim} \prod_{i \in I} \text{Hom}_R(N, M_i)$$

of abelian groups.

EXERCISE 1.18 (\star). Let K be a field. Let R be the set of ∞ by ∞ matrices

$$\begin{pmatrix} a_{11} & a_{12} & \cdots \\ a_{21} & a_{22} & \cdots \\ \vdots & \vdots & \end{pmatrix}$$

with $a_{ij} \in K$, and with the property that every column contains only finitely many non-zero elements. Verify that R (with the usual rule for matrix multiplication) is a ring. Show that the left R -modules R and $R \oplus R$ are isomorphic. Conclude that the condition that R is commutative cannot be dropped from Proposition 1.17.

CHAPTER 2

Exact sequences

Exact sequences form a useful and extensively used notational tool in algebra. They allow to replace tedious and verbose arguments involving kernels and quotients by quick and intuitive ‘diagram chases’.

1. Exact sequences

If $f: M_1 \rightarrow M_2$ and $g: M_2 \rightarrow M_3$ are R -module homomorphisms, then we say that the sequence

$$M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3$$

is *exact* if and only if the image of f is the kernel of g , as submodules of M_2 . For example: the sequence

$$0 \rightarrow M \rightarrow N$$

is exact if and only if $M \rightarrow N$ is injective, and the sequence

$$M \rightarrow N \rightarrow 0$$

is exact if and only if $M \rightarrow N$ is surjective.

A general sequence

$$\cdots \rightarrow M_{i-1} \xrightarrow{f_{i-1}} M_i \xrightarrow{f_i} M_{i+1} \rightarrow \cdots$$

is called exact if for every i we have $\ker f_i = \operatorname{im} f_{i-1}$ as submodules of M_i .

An exact sequence of the form

$$0 \rightarrow M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \rightarrow 0$$

is called a *short exact sequence*. Note that f induces an isomorphism

$$M_1 \cong \ker g$$

and g induces an isomorphism

$$\operatorname{coker} f \cong M_3.$$

We will often interpret the injective map f as the inclusion of a submodule M_1 into M_2 , and M_3 as the quotient of M_2 by the submodule M_1 , so that we can think of any short exact sequence as a sequence of the type

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_2/M_1 \longrightarrow 0.$$

2. The Five Lemma and the Snake Lemma

The Five Lemma and Snake Lemma are powerful and often-used lemmas about modules that are hard to state (and even harder to prove) without the language of commutative diagrams and exact sequences. The proofs are classic examples of ‘diagram chasing’. Such arguments are often fairly easy to verify by tracing elements around the diagram on the blackboard or a piece of paper, but are sometimes headache-provokingly resistant to being rendered or read in prose.

THEOREM 2.1 (Five Lemma). *Let R be a ring. Consider a commutative diagram of R -modules*

$$\begin{array}{ccccccccc} M_1 & \longrightarrow & M_2 & \longrightarrow & M_3 & \longrightarrow & M_4 & \longrightarrow & M_5 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow f_4 & & \downarrow f_5 \\ N_1 & \longrightarrow & N_2 & \longrightarrow & N_3 & \longrightarrow & N_4 & \longrightarrow & N_5 \end{array}$$

with exact rows. If f_1, f_2, f_4, f_5 are isomorphisms, then so is f_3 .

In fact, the proof will show that it suffices to assume that f_1 is surjective, f_5 is injective, and f_2 and f_4 are isomorphisms.

PROOF. The proof consists of two parts, one showing that f_3 is injective, the other that it is surjective. Both parts require only part of the hypotheses in the theorem.

Claim. If f_1 is surjective, and f_2 and f_4 are injective, then f_3 is injective.

Indeed, assume $f_3(x) = 0$ for some $x \in M_3$. We need to show that $x = 0$. Let $x' \in M_4$ be the image of x . By the commutativity of the diagram, $f_4(x') = 0$. But f_4 was injective, hence $x' = 0$. It follows that $x \in M_3$ is the image of some element $y \in M_2$. By commutativity, $f_2(y)$ maps to zero in N_3 , hence $f_2(y)$ is the image of some element $z \in N_1$. By the assumption on f_1 , there is a $\tilde{z} \in M_1$ with $f_1(\tilde{z}) = z$.

Consider the image y' of \tilde{z} in M_2 . By commutativity, we have $f_2(y') = f_2(y)$, but since f_2 is injective, this implies $y' = y$. We see that

$x \in M_3$ is the image of some element \tilde{z} in M_1 , and hence by exactness we conclude $x = 0$.

Claim. If f_5 is injective, and f_2 and f_4 are surjective, then f_3 is surjective. The proof of this second claim is left to the reader, see Exercise 2.3.

The theorem follows immediately from the above two claims. \square

THEOREM 2.2 (Snake Lemma). *Let R be a ring. Let*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & M_1 & \xrightarrow{\alpha} & M_2 & \xrightarrow{\beta} & M_3 & \longrightarrow & 0 \\ & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \\ 0 & \longrightarrow & N_1 & \xrightarrow{\alpha'} & N_2 & \xrightarrow{\beta'} & N_3 & \longrightarrow & 0 \end{array}$$

be a commutative diagram in which both horizontal rows are short exact sequences. Then there is a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \ker f_1 & \longrightarrow & \ker f_2 & \longrightarrow & \ker f_3 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & M_1 & \xrightarrow{\alpha} & M_2 & \xrightarrow{\beta} & M_3 & \longrightarrow & 0 \\ & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \\ 0 & \longrightarrow & N_1 & \xrightarrow{\alpha'} & N_2 & \xrightarrow{\beta'} & N_3 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ & & \text{coker } f_1 & \longrightarrow & \text{coker } f_2 & \longrightarrow & \text{coker } f_3 & \longrightarrow & 0 \end{array}$$

of R -modules in which the maps $\ker f_i \rightarrow M_i$ and $N_i \rightarrow \text{coker } f_i$ are the natural inclusions and projections, and in which the sequence

$$0 \rightarrow \ker f_1 \rightarrow \ker f_2 \rightarrow \ker f_3 \rightarrow \text{coker } f_1 \rightarrow \text{coker } f_2 \rightarrow \text{coker } f_3 \rightarrow 0$$

is exact.

SKETCH OF PROOF. We only give the most interesting part of the proof: the construction of the ‘snake’ map

$$d: \ker f_3 \rightarrow \text{coker } f_1.$$

Let $x \in \ker f_3 \subset M_3$. Since the map $\beta: M_2 \rightarrow M_3$ is surjective, there is a $y \in M_2$ with $\beta(y) = x$. By the commutativity of the right square, we have

$$\beta'(f_2(y)) = f_3(\beta(y)) = f_3(x) = 0.$$

So $f_2(y) \in \ker \beta' = \text{im } \alpha'$, hence there is a $z \in N_1$ with $\alpha'(z) = f_2(y)$. Note that z is unique, as the map α' is injective. We defined $d(x)$ as the element $\bar{z} \in \text{coker } f_1 = N_1/f_1(M_1)$.

We must check that this is well-defined, since our construction depended on the choice of $y \in M_2$ with $\beta(y) = x$. Let $y' \in M_2$ be another element with $\beta(y') = x$, leading to a $z' \in N_1$ as above. Since $\beta(y' - y) = x - x = 0$ there is a unique $\delta \in M_1$ with $y' - y = \alpha(\delta)$. Now the commutativity of the diagram shows $z' - z = f_1(\delta)$ in N_1 , and hence $\bar{z}' = \bar{z}$ in $N_1/f_1(M_1)$, as we had to show. \square

3. Split short exact sequences

Let M and N be R -modules. Then their direct sum fits into a short exact sequence

$$0 \longrightarrow M \xrightarrow{i} M \oplus N \xrightarrow{p} N \longrightarrow 0$$

where the maps are the natural inclusion $i: x \mapsto (x, 0)$, and projection $p: (x, y) \mapsto y$. It is often convenient to be able to recognize if a given short exact sequence is of the above special form.

THEOREM 2.3 (Splitting lemma). *Let*

$$0 \longrightarrow M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \longrightarrow 0$$

be an exact sequence of R -modules. Then the following are equivalent:

- (1) *there is a homomorphism $h: M_2 \rightarrow M_1$ such that $hf = \text{id}_{M_1}$*
- (2) *there is a homomorphism $s: M_3 \rightarrow M_2$ such that $gs = \text{id}_{M_3}$*
- (3) *there is an isomorphism $\varphi: M_2 \xrightarrow{\sim} M_1 \oplus M_3$ such that the diagram*

$$\begin{array}{ccccccc} 0 & \longrightarrow & M_1 & \xrightarrow{f} & M_2 & \xrightarrow{g} & M_3 & \longrightarrow & 0 \\ & & \downarrow \text{id} & & \downarrow \varphi & & \downarrow \text{id} & & \\ 0 & \longrightarrow & M_1 & \xrightarrow{i} & M_1 \oplus M_3 & \xrightarrow{p} & M_3 & \longrightarrow & 0 \end{array}$$

commutes (where $i(x) = (x, 0)$ and $p(x, y) = y$.)

If these conditions hold, we say that the sequence is *split* or *split exact*. The map h is called a *retraction* of f , and the map s a *section* of g . See Exercise 2.12 for examples of short exact sequences that are not split.

PROOF OF THEOREM 2.3. We first show that (3) implies (2). Indeed, the map

$$M_3 \rightarrow M_2, y \mapsto \varphi^{-1}(0, y)$$

is a section of g .

Next, we show that (2) implies (1), so assume that $s: M_3 \rightarrow M_2$ is a section. We will construct a retraction $h: M_2 \rightarrow M_1$. Let $x \in M_2$. Consider the element

$$y := x - s(g(x)) \in M_2.$$

Then, since $gs = \text{id}$ we have

$$g(y) = g(x) - g(s(g(x))) = g(x) - g(x) = 0.$$

So $y \in \ker g = \text{im } f$, and since f is injective, there is a unique $z \in M_1$ with $f(z) = y$. Define $h(x) := z$. One checks that h is indeed a retraction.

Finally, to show that (1) implies (3), assume that $h: M_2 \rightarrow M_1$ is a retraction. Then consider the map

$$\varphi: M_2 \rightarrow M_1 \oplus M_3, x \mapsto (h(x), g(x)).$$

Note that this map is an R -module homomorphism. We verify that it makes the diagram commute. We start with the left square. Take an $x \in M_1$ in the left-top corner of this square. Going down and then right, it gets mapped to $i(\text{id}(x)) = (x, 0) \in M_1 \oplus M_3$. Following the other path, we end up with $\varphi(f(x)) = (h(f(x)), g(f(x)))$. But now, $h(f(x)) = x$ because h is a retraction, and $g(f(x)) = 0$ because $\text{im } f = \ker g$ by the hypothesis that the sequence is exact. We conclude that $\varphi(f(x)) = (x, 0)$ and that the left square indeed commutes. For the right-hand square, take $x \in M_2$. Then one path yields $\text{id}(g(x)) = g(x)$, and following the other path, we obtain $p(\varphi(x)) = p(h(x), g(x)) = g(x)$. These agree, so we conclude that the diagram indeed commutes. Finally, by Exercise 2.5 we see that the map φ is automatically an isomorphism, which shows that (3) indeed follows from (1). \square

Exercises

EXERCISE 2.1. Show that

$$0 \longrightarrow M \longrightarrow 0$$

is exact if and only if M is the zero module.

EXERCISE 2.2. Let $f: M \rightarrow N$ be an R -module homomorphism. Show that there is an exact sequence

$$0 \longrightarrow \ker f \longrightarrow M \xrightarrow{f} N \longrightarrow \operatorname{coker} f \longrightarrow 0$$

of R -modules.

EXERCISE 2.3. Complete the proof of the Five Lemma (Theorem 2.1): show that if f_2 and f_4 are surjective, and if f_5 is injective, then f_3 is surjective.

EXERCISE 2.4. Let

$$\begin{array}{ccccccc} M_1 & \longrightarrow & M_2 & \longrightarrow & M_3 & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow & & \\ N_1 & \longrightarrow & N_2 & \longrightarrow & N_3 & \longrightarrow & 0 \end{array}$$

be a commutative diagram of R -modules with exact rows. Show that there exists a unique R -linear map $f_3: M_3 \rightarrow N_3$ making the resulting diagram commute.

EXERCISE 2.5. Consider a commutative diagram of R -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & M & \xrightarrow{\alpha} & E & \longrightarrow & N \longrightarrow 0 \\ & & \downarrow \operatorname{id}_M & & \downarrow f & & \downarrow \operatorname{id}_N \\ 0 & \longrightarrow & M & \longrightarrow & E' & \longrightarrow & N \longrightarrow 0 \end{array}$$

in which both rows are short exact sequences. Deduce from the snake or five lemma that f must be an isomorphism. Show that f is an isomorphism without using the snake or five lemma. (Hint for surjectivity: given $y \in E'$ choose an $x \in E$ with same image as y in N . Show that there is an $z \in M$ with $f(\alpha(z) + x) = y$.)

EXERCISE 2.6. Give an example of a diagram as in Theorem 2.2, for which the ‘snake map’ $d: \ker f_3 \rightarrow \operatorname{coker} f_1$ is non-zero.

EXERCISE 2.7. Let R be a ring and let

$$\begin{array}{ccc} M_1 & \hookrightarrow & M_2 \\ \downarrow \alpha_1 & & \downarrow \alpha_2 \\ N_1 & \hookrightarrow & N_2 \end{array}$$

be a commutative diagram of R -modules, in which the two horizontal maps are injective. Show that there exists an R -module E and an exact sequence

$$0 \longrightarrow \ker \alpha_1 \longrightarrow \ker \alpha_2 \longrightarrow E \longrightarrow \operatorname{coker} \alpha_1 \longrightarrow \operatorname{coker} \alpha_2$$

of R -modules.

EXERCISE 2.8. Let R be a ring, let

$$(2) \quad 0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3$$

be an exact sequence of R -modules, and let N be an R -module. Show that there is an exact sequence of abelian groups

$$0 \longrightarrow \operatorname{Hom}_R(N, M_1) \longrightarrow \operatorname{Hom}_R(N, M_2) \longrightarrow \operatorname{Hom}_R(N, M_3).$$

Give an example to show that the exactness of $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ need not imply that the map $\operatorname{Hom}_R(N, M_2) \rightarrow \operatorname{Hom}_R(N, M_3)$ is surjective.

EXERCISE 2.9. Let R be a ring, let

$$M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

be an exact sequence of R -modules, and let N be an R -module. Show that there is an exact sequence of abelian groups

$$0 \longrightarrow \operatorname{Hom}_R(M_3, N) \longrightarrow \operatorname{Hom}_R(M_2, N) \longrightarrow \operatorname{Hom}_R(M_1, N).$$

Give an example to show that the exactness of $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ need not imply that the map $\operatorname{Hom}_R(M_2, N) \rightarrow \operatorname{Hom}_R(M_1, N)$ is surjective.

EXERCISE 2.10. Let I and J be left ideals in a ring R . Show that there are exact sequences

$$0 \rightarrow I \cap J \rightarrow I \oplus J \rightarrow I + J \rightarrow 0$$

and

$$0 \rightarrow R/(I \cap J) \rightarrow R/I \oplus R/J \rightarrow R/(I + J) \rightarrow 0$$

of R -modules.

EXERCISE 2.11. Let R be a commutative ring and let

$$0 \longrightarrow M \longrightarrow E \longrightarrow N \longrightarrow 0$$

be a short exact sequence of R -modules. Let $I := \text{Ann}_R M$ and $J := \text{Ann}_R N$ (see Exercise 1.12). Show that

$$IJ \subset \text{Ann}_R E \subset I \cap J$$

as ideals in R .

EXERCISE 2.12. Show that the short exact sequences of \mathbf{Z} -modules

$$0 \longrightarrow \mathbf{Z} \xrightarrow{2} \mathbf{Z} \longrightarrow \mathbf{Z}/2\mathbf{Z} \longrightarrow 0$$

and

$$0 \longrightarrow \mathbf{Z}/2\mathbf{Z} \xrightarrow{2} \mathbf{Z}/4\mathbf{Z} \longrightarrow \mathbf{Z}/2\mathbf{Z} \longrightarrow 0$$

are *not* split. (Here the ‘2’ above the arrows are shorthand for the maps $x \mapsto 2x$ and $\bar{x} \mapsto 2\bar{x}$.)

EXERCISE 2.13. Let K be a field. Show that every short exact sequence of K -modules is split exact.

EXERCISE 2.14. Let G be a finite group, and K a field of characteristic zero. *Maschke’s theorem* asserts that for every representation V of G over K , and for every G -stable subspace $W \subset V$ there exists a G -stable complement $U \subset V$. Show that every short exact sequence of $K[G]$ -modules is split.

EXERCISE 2.15 (\star). Let G be the cyclic group of 2 elements. Consider the group ring $R := \mathbf{F}_2[G]$. Give an example of a non-split short exact sequence of R -modules. Show that not every representation of the group G over the field \mathbf{F}_2 is isomorphic to a direct sum of irreducible representations.

EXERCISE 2.16. Let K be a field. Consider the subring

$$R := \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid a, b, c \in K \right\}$$

of the ring $\text{Mat}(2, K)$ of two-by-two matrices over K . Let M be the module of column vectors $\begin{pmatrix} x \\ y \end{pmatrix}$ on which R acts by the usual matrix multiplication:

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax + by \\ cy \end{pmatrix}$$

Show that $N := \left\{ \begin{pmatrix} x \\ 0 \end{pmatrix} \mid x \in K \right\} \subset M$ is a sub- R -module, and that the short exact sequence of R -modules

$$0 \longrightarrow N \longrightarrow M \longrightarrow M/N \longrightarrow 0$$

does not split.

EXERCISE 2.17. Let R be a ring and let M and N be R -modules. Show that any short exact sequence of the form

$$0 \longrightarrow M \longrightarrow N \longrightarrow R^n \longrightarrow 0$$

is split.

EXERCISE 2.18. Let R be a ring and let $I \subset R$ be a left ideal. Show that a short exact sequence of left R -modules of the form

$$0 \longrightarrow M \longrightarrow N \xrightarrow{\pi} R/I \longrightarrow 0$$

splits if and only if there exists an $x \in N$ with $\pi(x) = 1 + I$ and $rx = 0$ for all $r \in I$.

EXERCISE 2.19. Let

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

be a split short exact sequence of R -modules, and let N be an R -module. Show that the induced sequences

$$0 \longrightarrow \text{Hom}_R(N, M_1) \longrightarrow \text{Hom}_R(N, M_2) \longrightarrow \text{Hom}_R(N, M_3) \longrightarrow 0$$

and

$$0 \longrightarrow \text{Hom}_R(M_3, N) \longrightarrow \text{Hom}_R(M_2, N) \longrightarrow \text{Hom}_R(M_1, N) \longrightarrow 0$$

are exact.

EXERCISE 2.20. Let R be a ring and let

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

be a short exact sequence of R -modules. Show that if M_1 is free of rank n_1 and M_3 is free of rank n_3 , then M_2 is free of rank $n_1 + n_3$.

CHAPTER 3

Finitely generated modules over a PID

1. Introduction

The classification of finite abelian groups states that for every finite abelian group A there are prime numbers p_i (not necessarily distinct) and exponents $e_i \geq 1$ such that

$$A \cong (\mathbf{Z}/p_1^{e_1}\mathbf{Z}) \times \cdots \times (\mathbf{Z}/p_n^{e_n}\mathbf{Z}).$$

The existence of Jordan normal forms states that for every square matrix P over \mathbf{C} there exist complex numbers λ_i (not necessarily distinct) and integers $e_i \geq 1$ so that P is conjugate to a block diagonal matrix with blocks

$$\begin{pmatrix} \lambda_i & 0 & \cdots & 0 & 0 \\ 1 & \lambda_i & \cdots & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda_i & 0 \\ 0 & 0 & \cdots & 1 & \lambda_i \end{pmatrix}$$

of size e_i .

In this chapter, we will see that these two theorems are just two instances of one and the same theorem about finitely generated modules over a principal ideal domain. In the first case, the PID will be \mathbf{Z} , in the second case it will be $\mathbf{C}[X]$.

2. Review of principal ideal domains

Let R be a PID (principal ideal domain). Recall that this means that R is a domain (a nonzero commutative ring without zero divisors), and that for every ideal $I \subset R$ there is an $r \in R$ with $I = (r) = Rr$. For our purposes, the most important examples are $R = \mathbf{Z}$ and $R = K[X]$ with K a field. Other important examples are the ring of power series $K[[X]]$, the ring of p -adic integers \mathbf{Z}_p , and the ring of Gaussian integers $\mathbf{Z}[i]$. Also a field K is a PID, but of a rather trivial kind.

PID's are unique factorization domains. This means that every non-zero element r in a principal ideal domain R can be written as

$$r = up_1^{e_1} \cdots p_n^{e_n}$$

with $u \in R^\times$, with the $p_i \in R$ irreducible, and with e_i non-negative integers. Moreover, such factorization is unique up to multiplying u and the p_i 's by units, and up to permuting the factors.

Using this prime factorization, we can define greatest common divisors $\gcd(r, s)$ of two non-zero elements of R . They are uniquely determined up to units (for example both 2 and -2 are a gcd of 4 and 6 in \mathbf{Z}). Similarly, we can define gcds of any sequence r_1, \dots, r_n of elements of R which is not identically zero (ignoring the zeroes in the sequence).

An element $d \in R$ is a gcd of r_1, \dots, r_n if and only if d generates the ideal (r_1, \dots, r_n) of R . In particular, there are a_1, \dots, a_n in R with $d = a_1r_1 + \cdots + a_nr_n$. For example, if r and s are coprime (have no common prime factor), then there are a and b in R with $ar + bs = 1$.

3. Free modules of finite rank over a PID

PROPOSITION 3.1. *Let R be a PID and $M \subset R^n$ a submodule. Then $M \cong R^k$ for some $k \leq n$.*

Hence over a PID a submodule of a free module of finite rank is itself free of finite rank. The condition that R be a PID cannot be dropped from the proposition, see Exercise 3.3.

PROOF OF PROPOSITION 3.1. We use induction on n . For $n = 0$ we have $M = R^n = 0$, and M is indeed free of rank 0. Assume that the proposition has been shown to hold for submodules of R^{n-1} . Consider the projection

$$\pi: R^n \rightarrow R, (r_1, \dots, r_n) \rightarrow r_n$$

with kernel $R^{n-1} \times \{0\}$. Then we have a short exact sequence

$$0 \rightarrow M \cap \ker \pi \rightarrow M \rightarrow \pi(M) \rightarrow 0.$$

By the induction hypothesis, we have that the submodule $M \cap \ker \pi$ of $R^{n-1} \times \{0\}$ is free of rank $k \leq n - 1$. If $\pi(M) = 0$, then we are done. If $\pi(M) \neq 0$, then $\pi(M)$ is an ideal in R , hence a principal ideal, hence $\pi(M) \cong R$ as R -module. By Exercise 2.17 any short exact sequence of R -modules of the form

$$0 \rightarrow N \rightarrow M \rightarrow R \rightarrow 0$$

splits, and we find $M \cong R \oplus (M \cap \ker \pi) \cong R^{k+1}$ with $k+1 \leq n$. \square

COROLLARY 3.2. *Let R be a PID and M a finitely generated R -module. Then there exists an exact sequence*

$$0 \longrightarrow F_1 \longrightarrow F_2 \longrightarrow M \longrightarrow 0$$

with F_1 and F_2 free R -modules of finite rank.

PROOF. Since M is finitely generated, by Proposition 1.13 there is a surjection $F_2 \rightarrow M$ with F_2 a free module of finite rank. The kernel $F_1 \subset F_2$ is also free of finite rank, thanks to Proposition 3.1. \square

4. Structure of finitely generated modules over a PID

The main theorem of this chapter is the following structure theorem.

THEOREM 3.3. *Let R be a principal ideal domain and M a finitely generated R -module. Then there exists an integer n and non-zero ideals I_1, \dots, I_k of R such that*

$$M \cong R^n \oplus R/I_1 \oplus \cdots \oplus R/I_k$$

as R -modules.

For the proof we need the notion of *content* of an element of a module. Let R be a commutative ring and let M be a an R -module. An element $x \in M$ determines a map

$$(3) \quad \text{Hom}_R(M, R) \rightarrow R, f \mapsto f(x).$$

This map is R -linear, hence the image is an ideal. We denote it by $c_M(x)$, and call it the *content* of x .

LEMMA 3.4. *If M is free of finite rank, and $x \in M$ is non-zero, then $c_M(x)$ is a non-zero ideal in R .*

PROOF. Without loss of generality, we may assume $M = R^n$ and $x = (x_1, \dots, x_n)$. Since x is non-zero, there exists an i with $x_i \neq 0$. Consider the map

$$\text{pr}_i: R^n \rightarrow R, (y_1, \dots, y_n) \mapsto y_i.$$

We have $\text{pr}_i \in \text{Hom}_R(M, R)$ and $\text{pr}_i(x) \neq 0$, hence $c_M(x) \neq 0$. \square

Note that the coordinates x_i of $x \in M \cong R^n$ depend on the choice of basis of M , but that the content ideal $c_M(x)$ is independent of such choice!

LEMMA 3.5. *Let R be a principal ideal domain, M a free R -module of finite rank, and $x \in M$ a non-zero element. Then there is a surjective R -linear map $f: M \rightarrow R$ such that $f(x) \in R$ generates the ideal $c_M(x) \subset R$.*

PROOF. Since R is a principal ideal domain, there exists an $f \in \text{Hom}_R(M, R)$ such that $f(x)$ generates the ideal $c_M(x)$. Moreover, by Lemma 3.4 we have $f(x) \neq 0$.

Consider the image of $f: M \rightarrow R$. This is an ideal $I \subset R$, and it suffices to show that $I = R$. As R is a PID, the ideal I is generated by some element $r \in R$. For every $y \in M$ we have $f(y) \in I$ and (since R is a domain) there exists a unique $g(y) \in R$ such that

$$f(y) = g(y) \cdot r.$$

Now consider the element $g(x) \in R$. By definition of $c_M(x)$ we have $g(x) \in c_M(x)$. Since $f(x)$ generates $c_M(x)$ there is an $s \in R$ with $g(x) = sf(x)$. But we also have $f(x) = rg(x)$, hence $rs = 1$, hence $I = R$ and f is surjective. \square

PROOF OF THEOREM 3.3. By Corollary 3.2 every finitely generated R -module M sits in a short exact sequence

$$0 \longrightarrow F_1 \longrightarrow F_2 \longrightarrow M \longrightarrow 0$$

with the F_i free R -modules of finite rank. We will interpret F_1 as a submodule of F_2 .

The proof goes by induction on the rank of F_1 . If F_1 is zero, then $M \cong F_2 \cong R^n$ for some n , and the theorem holds.

Otherwise, let $x \in F_1$ be a non-zero element whose content $I := c_{F_2}(x)$ with respect to F_2 is maximal (amongst all the ideals in R of the form $c_{F_2}(x)$ with $x \in F_1$).

By Lemma 3.5 there is a surjective map $f: F_2 \rightarrow R$ such that $f(x)$ generates I .

Claim. $f(F_1) = I \subset R$. Indeed, since $f(x)$ generates I we certainly have $f(F_1) \supset I$. Conversely, for $z \in F_1$, let d be a gcd of $f(z)$ and $f(x)$. Then there exists $r, s \in R$ with $rf(z) + sf(x) = d$, and hence $f(rz + sx) = d$. By the maximality of the content of x we have that d must equal $f(x)$ up to a unit, hence $f(z)$ must be divisible by $f(x)$ and hence $f(z) \in I$.

Denote the kernel of $f: F_2 \rightarrow R$ by F'_2 (note that by Proposition 3.1, this is also a free module). We have a short exact sequence

$$0 \longrightarrow F'_2 \longrightarrow F_2 \xrightarrow{f} R \longrightarrow 0.$$

Similarly, let F'_1 be the kernel of the restriction $f: F_1 \rightarrow R$. We find a short exact ‘sub-sequence’

$$0 \longrightarrow F'_1 \longrightarrow F_1 \xrightarrow{f} I \longrightarrow 0.$$

The first sequence splits by the section $R \rightarrow F_2$, $1 \mapsto x/f(x)$, and this section restricts to a section $I \rightarrow F_1$ in the second short exact sequence.

We find $M \cong M' \oplus R/I$ with M' given by the short exact sequence

$$0 \longrightarrow F'_1 \longrightarrow F'_2 \longrightarrow M' \longrightarrow 0.$$

Since F'_2 has lower rank, the induction hypothesis guarantees

$$M' \cong R^{n'} \oplus R/I_1 \oplus \cdots \oplus R/I_k,$$

which finishes the proof. \square

5. Application to Jordan normal form

Theorem 3.3 has the following corollary.

COROLLARY 3.6. *Let R be a PID and M a finitely generated R -module. Then there exists an integer n , irreducible elements p_1, \dots, p_k of R and positive integers e_1, \dots, e_k such that*

$$M \cong R^n \oplus R/p_1^{e_1} R \oplus \cdots \oplus R/p_k^{e_k} R$$

as R -modules.

PROOF. By Theorem 3.3 it suffices to show that for every non-zero ideal I the R -module R/I can be written in the desired form. Let x be a generator of I , and consider its prime factorization

$$x = up_1^{e_1} \cdots p_k^{e_k}$$

with p_i pairwise non-associated primes. Then by the Chinese Remainder Theorem, we have

$$R/I \cong R/p_1^{e_1} R \oplus \cdots \oplus R/p_k^{e_k} R,$$

as we had to show. \square

We now consider two special cases of this corollary. The first one is a structure theorem for finitely generated and finite abelian groups.

THEOREM 3.7 (Classification of finitely generated abelian groups). *Let A be a finitely generated abelian group. Then there exists an integer n , prime numbers p_1, \dots, p_k , and positive integers e_1, \dots, e_k such that*

$$A \cong \mathbf{Z}^n \times \mathbf{Z}/p_1^{e_1}\mathbf{Z} \times \cdots \times \mathbf{Z}/p_k^{e_k}\mathbf{Z}.$$

If A is a finite abelian group then

$$A \cong \mathbf{Z}/p_1^{e_1}\mathbf{Z} \times \cdots \times \mathbf{Z}/p_k^{e_k}\mathbf{Z}.$$

PROOF. Apply Corollary 3.6 to the case $R = \mathbf{Z}$. \square

The second special case is a structure theorem for endomorphisms of finite-dimensional vector spaces over \mathbf{C} (or over an algebraically closed field).

THEOREM 3.8. *Let K be an algebraically closed field. Let V be a finite-dimensional vector space over K . Let $\alpha: V \rightarrow V$ be an endomorphism. Then there exist $\lambda_1, \dots, \lambda_k$ in K , positive integers e_1, \dots, e_k , and a decomposition*

$$V = V_1 \oplus \cdots \oplus V_k$$

such that $\alpha(V_i) \subset V_i$, and such that each V_i has a basis on which α is expressed as the standard Jordan matrix

$$\begin{pmatrix} \lambda_i & 0 & \cdots & 0 & 0 \\ 1 & \lambda_i & \cdots & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda_i & 0 \\ 0 & 0 & \cdots & 1 & \lambda_i \end{pmatrix}$$

of size e_i .

PROOF. As in Example 1.9, we turn the vector space V into a $K[X]$ -module, with X acting via the endomorphism α . Since V is finite-dimensional, it is finitely generated as a K -module, hence a fortiori also as a $K[X]$ -module.

Since K is algebraically closed, the irreducible elements of $K[X]$ are (up to units) the linear polynomials $X - \lambda$ with $\lambda \in K$. By Corollary 3.6, we have

$$V \cong K[X]^n \oplus \frac{K[X]}{(X - \lambda_1)^{e_1} K[X]} \oplus \cdots \oplus \frac{K[X]}{(X - \lambda_k)^{e_k} K[X]}.$$

Note that $K[X]$ is infinite-dimensional as a K -vector space, so we necessarily must have $n = 0$.

Without loss of generality, we may assume

$$V = K[X]/(X - \lambda)^e K[X]$$

for some $\lambda \in K$ and $e > 0$. Consider the elements

$$v_j := (\bar{X} - \lambda)^j \in V \quad j \in \{0, \dots, e-1\}.$$

Note that the v_j form a K -basis of V . The action of α on this basis is given by

$$\alpha(v_j) = X \cdot (\bar{X} - \lambda)^j = (\bar{X} - \lambda)^{j+1} + \lambda(\bar{X} - \lambda)^j$$

hence

$$\alpha(v_j) = \begin{cases} v_{j+1} + \lambda v_j & j < e-1 \\ \lambda v_j & j = e-1. \end{cases}$$

and we see that the matrix of α with respect to this basis is the standard Jordan block of eigenvalue λ and size e . \square

Exercises

EXERCISE 3.1. Let R be a commutative ring and $\mathfrak{m} \subset R$ a maximal ideal. Let M be an R -module. Show that $M/\mathfrak{m}M$ is a vector space over R/\mathfrak{m} . Show that if M is generated by x_1, \dots, x_n then $M/\mathfrak{m}M$ has dimension at most n over R/\mathfrak{m} .

EXERCISE 3.2. Let R be a commutative ring and M an R -module. An element $x \in M$ is called a *torsion* element if there exists a non-zero $r \in R$ with $rx = 0$.

Assume that R is a domain. Show that the torsion elements of an R -module M form a submodule of M .

Give an example to show that the condition that R is a domain cannot be dropped.

EXERCISE 3.3. Let R be an integral domain. Show that the following are equivalent:

- (1) every submodule of a free module of finite rank is free of finite rank,
- (2) R is a principal ideal domain.

EXERCISE 3.4. Let R be a PID and let M be an R -module which can be generated by n elements. Show that every submodule $N \subset M$ can be generated by n elements. Show that the condition that R is a PID cannot be dropped.

EXERCISE 3.5. Let R be a PID and let M be a finitely generated R -module such that for all $r \in R$ and $x \in M$ we have that $rx = 0$ implies $r = 0$ or $x = 0$. Show that M is free of finite rank.

EXERCISE 3.6. Let R be a ring and let

$$0 \longrightarrow M \longrightarrow E \longrightarrow N \longrightarrow 0$$

be a short exact sequence of R -modules. Assume that M can be generated by m elements, and that N can be generated by n elements. Show that E can be generated by $m + n$ elements.

EXERCISE 3.7. Let R be a PID, and let p_1 and p_2 be irreducible elements with $(p_1) \neq (p_2)$. Let e_1, e_2 be non-negative integers. Show that the only R -module homomorphism

$$R/p_1^{e_1}R \rightarrow R/p_2^{e_2}R$$

is the zero homomorphism.

EXERCISE 3.8. Let R be a PID and let $p \in R$ be irreducible. Let e_1, e_2 be non-negative integers. Show that there is an isomorphism of R -modules

$$\mathrm{Hom}_R(R/p^{e_1}R, R/p^{e_2}R) \cong R/p^eR$$

with $e = \min(e_1, e_2)$.

EXERCISE 3.9. Describe all $\mathbf{Z}[i]$ -modules with at most 10 elements, up to isomorphism.

EXERCISE 3.10. Let R be a PID and let $p \in R$ be an irreducible element. Let E be an R -module contained in a short exact sequence

$$0 \longrightarrow R/pR \xrightarrow{f} E \xrightarrow{g} R/pR \longrightarrow 0.$$

Show that either $E \cong R/pR \oplus R/pR$ or $E \cong R/p^2R$, that both options occur, and that $R/pR \oplus R/pR \not\cong R/p^2R$.

EXERCISE 3.11 (\star). Let

$$0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow A_3 \longrightarrow \cdots \longrightarrow A_n \longrightarrow 0$$

be an exact sequence of finite abelian groups. Show that the equality

$$\prod_i |A_i|^{(-1)^i} = 1$$

holds.

EXERCISE 3.12. Let K be a field and let $f_1, \dots, f_n \in K[X]$ be monic polynomials. Consider the $K[X]$ -module

$$V := K[X]/(f_1) \oplus \cdots \oplus K[X]/(f_n).$$

Show that the characteristic polynomial of the endomorphism $v \mapsto X \cdot v$ of the K -vector space V equals $\prod_i f_i$.

EXERCISE 3.13. Let K be a field and let V be a finite-dimensional vector space over K . Let α be an endomorphism of V . Assume that the characteristic polynomial of α is irreducible. Show that there is no proper non-zero subspace $W \subset V$ with $\alpha(W) \subset W$.

EXERCISE 3.14. Let K be a field and let V be a finite-dimensional vector space over K . Let α be an endomorphism of V . Assume that the characteristic polynomial of α is separable. Show that there are subspaces W_1, \dots, W_n of V , such that the following hold:

$$(1) \quad V = W_1 \oplus W_2 \oplus \cdots \oplus W_n,$$

- (2) $\alpha(W_i) \subset W_i$ for every i ,
- (3) the characteristic polynomial of $\alpha|_{W_i}$ is irreducible for every i .

Give an example to show that the condition that the characteristic polynomial is separable cannot be dropped.

EXERCISE 3.15 ($\star\star$). Let R be a PID. An R -module M is called *torsion* if for every $x \in M$ there is a non-zero $r \in R$ with $rx = 0$. For a finitely generated torsion R -module

$$M \cong R/I_1 \oplus \cdots \oplus R/I_k$$

denote by $F(M)$ the ideal $I_1 I_2 \cdots I_k$. Show that $F(M)$ is independent of the chosen decomposition. Show that if

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

is a short exact sequence of R -modules, and if M_1 and M_3 are finitely generated torsion R -modules, then M_2 is a finitely generated torsion R -module and $F(M_2) = F(M_1)F(M_3)$.

CHAPTER 4

Categories

1. Definition

DEFINITION 4.1. A *category* \mathcal{C} consists of the data of

- (1) a class of *objects* $\text{ob } \mathcal{C}$,
- (2) for every $X, Y \in \text{ob } \mathcal{C}$ a class $\text{Hom}_{\mathcal{C}}(X, Y)$
- (3) for every $X \in \text{ob } \mathcal{C}$ an element $\text{id}_X \in \text{Hom}_{\mathcal{C}}(X, X)$
- (4) for every $X, Y, Z \in \text{ob } \mathcal{C}$ a map

$$\text{Hom}_{\mathcal{C}}(X, Y) \times \text{Hom}_{\mathcal{C}}(Y, Z) \rightarrow \text{Hom}_{\mathcal{C}}(X, Z), (f, g) \mapsto gf,$$

called *composition*, subject to the conditions

- (C1) for every $X, Y, Z, T \in \text{ob } \mathcal{C}$ and for every $f \in \text{Hom}_{\mathcal{C}}(X, Y)$, $g \in \text{Hom}_{\mathcal{C}}(Y, Z)$ and $h \in \text{Hom}_{\mathcal{C}}(Z, T)$ the identity $h(gf) = (hg)f$ holds
- (C2) for every $X, Y \in \mathcal{C}$ and for every $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ the identities $f\text{id}_X = f$ and $\text{id}_Y f = f$ hold.

We often write $f: X \rightarrow Y$ instead of $f \in \text{Hom}_{\mathcal{C}}(X, Y)$, and think of f as being a ‘map from X to Y ’. However, the objects X and Y need not be sets, and expressions such as ‘ $x \in X$ ’ or ‘ $f(x) \in Y$ ’ can be completely meaningless.

REMARK 4.2. The use of the word ‘class’ is to avoid set-theoretical problems. We want to consider examples such as the category of all sets, but must be careful to avoid paradoxes. The class of all sets does not form a set itself, otherwise one could consider the subset of all the sets that are not contained in itself, leading to Russell’s paradox.

This subtlety in the definition of a category is mostly harmless, and in almost all applications of categories one can safely pretend the class of objects form a set. In fact, one can often restrict the objects to a suitable chosen sub-*set* of the given class without losing much.

2. Big examples

The notion of a category is modeled on the properties of the collection of all objects of a certain kind (sets, rings, spaces) together with the collection of all structure-preserving maps (functions, ring homomorphisms, continuous maps) between them. The most important examples are of this kind.

EXAMPLE 4.3 (The category of sets). The category **Set** with $\text{ob } \mathbf{Set}$ the class of all sets, $\text{Hom}_{\mathbf{Set}}(X, Y)$ the set of all maps from X to Y , id_X the identity map and the usual composition $gf := g \circ f$ forms a category.

EXAMPLE 4.4 (The category of topological spaces). The category **Top** with $\text{ob } \mathbf{Top}$ the class of all topological spaces, $\text{Hom}_{\mathbf{Top}}(X, Y)$ the set of continuous maps from X to Y and the usual identity and composition form a category. (Note that the composition of two continuous maps is continuous!).

EXAMPLE 4.5 (The categories of left and right R -modules). If R is a ring, we denote by ${}_R\mathbf{Mod}$ the category whose objects are the left R -modules, and whose morphisms are the R -module homomorphisms. So for $M, N \in \text{ob } {}_R\mathbf{Mod}$ we have

$$\text{Hom}_{{}_R\mathbf{Mod}}(M, N) := \text{Hom}_R(M, N).$$

Similarly, we denote by \mathbf{Mod}_R the category of right R -modules.

In the same style as the above examples, we have the category **Ring** of rings and ring homomorphisms, the category **CRing** of commutative rings and ring homomorphisms, the category **Grp** of groups and group homomorphisms, the category **Ab** of abelian groups and group homomorphisms, etcetera. Note that in all these examples the objects of the categories are sets equipped with some extra structure, and the morphisms are functions that are compatible with the structure. This is the case for many, but certainly not all, commonly used categories.

3. Small examples

A category is also a mathematical object in its own right, and one can write down explicit examples by specifying the objects and maps, in the same way one can specify say a ring by giving its elements, the addition, and multiplication.

EXAMPLE 4.6 (One arrow). Consider the category \mathcal{C} consisting of precisely two objects, X and Y , and with precisely three maps: id_X , id_Y , and a map $f: X \rightarrow Y$. We can render this category in a picture:

$$\text{id}_X \circlearrowleft \bullet \xrightarrow{f} \bullet \circlearrowright \text{id}_Y \\ X \qquad Y$$

EXAMPLE 4.7 (A group as a category). Let G be a group. Consider the category BG with one object \star (so $\text{ob BG} = \{\star\}$), and with $\text{Hom}_{\text{BG}}(\star, \star) := G$, $\text{id}_\star := 1 \in G$, and where composition is defined by multiplication in G . Note that axiom (C1) follows from the associativity of the group operation, and axiom (C2) from the axiom for the neutral element $1 \in G$.

EXAMPLE 4.8 (Discrete category). If S is a set, then S defines a category \mathcal{C} with $\text{ob } \mathcal{C} := S$ and with

$$\text{Hom}_{\mathcal{C}}(x, y) := \begin{cases} \{\text{id}\} & \text{if } x = y \\ \emptyset & \text{if } x \neq y \end{cases}$$

for all $x, y \in S$.

EXAMPLE 4.9 (Pre-ordered set). A *pre-ordered set* is a set S equipped with a relation \leq that is reflexive ($x \leq x$) and transitive ($x \leq y$ and $y \leq z$ implies $x \leq z$). A pre-ordered set defines a category \mathcal{C} with $\mathcal{C} := S$ and with

$$\text{Hom}_{\mathcal{C}}(x, y) := \begin{cases} \{\star\} & \text{if } x \leq y \\ \emptyset & \text{if } x \not\leq y \end{cases}$$

where $\{\star\}$ denotes any singleton.

EXAMPLE 4.10 (The category of matrices). Let R be a ring. Then we can form a category \mathcal{C} with $\text{ob } \mathcal{C} := \{0, 1, 2, \dots\}$, with

$$\text{Hom}_{\mathcal{C}}(n, m) := \text{Mat}_{m,n}(R) := \{m \times n \text{ matrices over } R\},$$

and with composition being matrix multiplication

$$\text{Mat}_{\ell,k} \times \text{Mat}_{m,\ell} \rightarrow \text{Mat}_{m,k}, (A, B) \mapsto BA.$$

The identity element $\text{id}_n: n \rightarrow n$ is the $n \times n$ identity matrix. Axiom (C1) corresponds to the associativity of matrix multiplication.

DEFINITION 4.11. A category is called *locally small* if for every X and Y in $\text{ob } \mathcal{C}$ the class $\text{Hom}_{\mathcal{C}}(X, Y)$ is a set. It is called *small* if moreover the class of objects $\text{ob } \mathcal{C}$ is itself a set.

EXAMPLES 4.12. All the examples in sections 2 and 3 are locally small. The examples in 3 are in fact small. However, the examples in 2 are not small, essentially because the class of all sets is not a set.

Finally, we discuss two ways of constructing new categories out of old ones. The first is the opposite category, which is obtained by formally reversing all the arrows in a given category:

DEFINITION 4.13. The *opposite* or *dual* of a category \mathcal{C} is the category \mathcal{C}^{opp} with $\text{ob } \mathcal{C}^{\text{opp}} := \text{ob } \mathcal{C}$ and with

$$\text{Hom}_{\mathcal{C}^{\text{opp}}}(X, Y) := \text{Hom}_{\mathcal{C}}(Y, X).$$

Composition is done ‘the other way around’:

$$\text{Hom}_{\mathcal{C}^{\text{opp}}}(X, Y) \times \text{Hom}_{\mathcal{C}^{\text{opp}}}(Y, Z) \rightarrow \text{Hom}_{\mathcal{C}^{\text{opp}}}(X, Z), (f, g) \mapsto fg$$

where $f: Y \rightarrow X$ and $g: Z \rightarrow Y$ and $fg: Z \rightarrow X$ are maps in \mathcal{C} .

DEFINITION 4.14. If \mathcal{C} and \mathcal{D} are categories, then the *product category* $\mathcal{C} \times \mathcal{D}$ is the category whose objects are pairs (X, Y) with $X \in \text{ob } \mathcal{C}$ and $Y \in \text{ob } \mathcal{D}$, and whose morphisms are pairs of morphisms:

$$\text{Hom}_{\mathcal{C} \times \mathcal{D}}((X, Y), (X', Y')) := \text{Hom}_{\mathcal{C}}(X, X') \times \text{Hom}_{\mathcal{D}}(Y, Y').$$

4. Isomorphisms, monomorphisms, epimorphisms

In general it makes no sense to ask if a morphism $f: X \rightarrow Y$ in a category \mathcal{C} is bijective, injective, or surjective. The objects X and Y are just elements of some class $\text{ob } \mathcal{C}$, and the morphism f is just an element of some class $\text{Hom}_{\mathcal{C}}(X, Y)$. It makes no sense to talk about elements of X or Y .

However, there are special classes of morphisms that in many ways behave like bijective, injective, or surjective maps.

DEFINITION 4.15. Let $f: X \rightarrow Y$ be a morphism in a category \mathcal{C} . We say that f is an *isomorphism* if there exists a morphism $g: Y \rightarrow X$ such that $fg = \text{id}_Y$ and $gf = \text{id}_X$.

EXAMPLE 4.16. An isomorphism in **Set** is a bijection. An isomorphism in **Grp** is a group isomorphism. An isomorphism in **Top** is

a homeomorphism (note that this is stronger than being a bijective continuous map!). An isomorphism in the category of matrices (Example 4.10) is an invertible square matrix.

DEFINITION 4.17. A morphism $f: X \rightarrow Y$ in a category \mathcal{C} is called a *monomorphism* if for every object T and for every pair of maps $g_1, g_2: T \rightarrow X$ with $fg_1 = fg_2$ we have $g_1 = g_2$.

EXAMPLE 4.18. We claim that the monomorphisms in **Set** are precisely the injective functions. Indeed, if f is injective, then clearly $fg_1 = fg_2$ implies $g_1 = g_2$. Conversely, let $f: X \rightarrow Y$ be a monomorphism. Let $x_1, x_2 \in X$ with $f(x_1) = f(x_2)$. Consider the one-element set $T = \{\star\}$ and the maps $g_i: T \rightarrow X$, $\star \mapsto x_i$. Then $fg_1 = fg_2$, hence $g_1 = g_2$, and therefore $x_1 = x_2$.

EXAMPLE 4.19. By a similar argument one shows that the monomorphisms in **Grp** are precisely the injective group homomorphisms, see Exercise 4.5.

DEFINITION 4.20. A morphism $f: X \rightarrow Y$ in a category \mathcal{C} is called an *epimorphism* if for every $T \in \text{ob } \mathcal{C}$ and for every pair of maps g_1, g_2 from Y to T with $g_1f = g_2f$ we have $g_1 = g_2$.

EXAMPLE 4.21. We claim that the epimorphisms in **Set** are precisely the surjective functions. Indeed, if f is surjective and $g_1f = g_2f$, then certainly $g_1 = g_2$. Conversely, let $f: X \rightarrow Y$ be an epimorphism. Assume that f is not surjective, and pick an element $z \in Y$ not in the image of f . Consider the set $T := \{0, 1\}$ and maps

$$g_1: Y \rightarrow T, y \mapsto 0$$

and

$$g_2: Y \rightarrow T, y \mapsto \begin{cases} 0 & y \neq z \\ 1 & y = z \end{cases}$$

Then $g_1f = g_2f$, but $g_1 \neq g_2$, a contradiction.

5. Initial and final objects

Some categories have special objects, called initial (or cofinal) and final objects.

DEFINITION 4.22. Let \mathcal{C} be a category. An object $X \in \text{ob } \mathcal{C}$ is called *final* in \mathcal{C} if for every $Y \in \text{ob } \mathcal{C}$ there is a unique morphism $Y \rightarrow X$.

DEFINITION 4.23. Let \mathcal{C} be a category. An object $X \in \text{ob } \mathcal{C}$ is called *initial* or *cofinal* in \mathcal{C} if for every $Y \in \text{ob } \mathcal{C}$ there is a unique morphism $X \rightarrow Y$.

EXAMPLES 4.24. In **Top** the empty space \emptyset is an initial object, and a one-point space $\{\star\}$ is final. In ${}_R\mathbf{Mod}$ the zero module 0 is both initial and final. In **Ring** the ring \mathbf{Z} is initial and the zero ring 0 is final. In **Grp** the trivial group $\{1\}$ is both initial and final.

PROPOSITION 4.25. *If it exists, a final object in a category \mathcal{C} is unique up to unique isomorphism.*

PROOF. Assume X_1 and X_2 are final. We need to show that there exists a unique isomorphism $f: X_1 \rightarrow X_2$.

Since X_2 is final, there exists a map $f: X_1 \rightarrow X_2$. Since X_1 is final, there exists a map $g: X_2 \rightarrow X_1$. Consider the composite $fg: X_2 \rightarrow X_2$. Since X_2 is final, there is a unique map $X_2 \rightarrow X_2$, so $fg = \text{id}_{X_2}$. Similarly $gf = \text{id}_{X_1}$. So we see that f is an isomorphism $X_1 \rightarrow X_2$.

Now assume that there are *two* isomorphisms $f_1, f_2: X_1 \rightarrow X_2$. Then again, because X_2 is final, we must have $f_1 = f_2$, which shows that f is unique. \square

Like so many statements about categories, this proposition has a co-proposition:

PROPOSITION 4.26. *If it exists, an initial object in a category \mathcal{C} is unique up to unique isomorphism.*

PROOF. Reverse the arrows in the proof of Proposition 4.25. \square

Exercises

EXERCISE 4.1 (Automorphism group of an object). Let \mathcal{C} be a (locally small) category and X an object in \mathcal{C} . Show that

$$\text{Aut}_{\mathcal{C}}(X) := \{f: X \rightarrow X \mid f \text{ is an isomorphism}\}$$

forms a group under composition.

EXERCISE 4.2. Let \mathcal{C} be a (locally small) category and X and Y objects in \mathcal{C} . Assume that X and Y are isomorphic. Show that $\text{Aut}_{\mathcal{C}}(X)$ and $\text{Aut}_{\mathcal{C}}(Y)$ are isomorphic groups.

EXERCISE 4.3. Let $f: X \rightarrow Y$ be morphism in a category \mathcal{C} . Show that f has at most one two-sided inverse.

EXERCISE 4.4. Let \mathcal{C} be a category. Define $\text{ob } \mathcal{C}^{\times} := \text{ob } \mathcal{C}$ and

$$\text{Hom}_{\mathcal{C}^{\times}}(X, Y) := \{f \in \text{Hom}_{\mathcal{C}}(X, Y) \mid f \text{ is an isomorphism}\}.$$

Show that \mathcal{C}^{\times} (with composition and identity maps inherited from \mathcal{C}) is a category.

EXERCISE 4.5. Show that the monomorphisms in \mathbf{Grp} are precisely the injective group homomorphisms. (Hint: given $x \in G$ there is a group homomorphism $\mathbf{Z} \rightarrow G$ which maps 1 to x).

EXERCISE 4.6. Let R be a ring. Show that the monomorphisms in ${}_R\mathbf{Mod}$ are precisely the injective R -module homomorphisms.

EXERCISE 4.7. Show that the epimorphisms in ${}_R\mathbf{Mod}$ are precisely the surjective homomorphisms. (Hint: given $f: M \rightarrow N$ non-surjective, consider the module $\text{coker } f$.)

EXERCISE 4.8. Show that the map $\mathbf{Z} \rightarrow \mathbf{Q}$ in the category of rings is an epimorphism. Conclude that there exist morphisms that are both mono and epi, but not isomorphisms.

EXERCISE 4.9. Let \mathbf{Haus} be the category of Hausdorff topological spaces and continuous maps. Let $f: X \rightarrow Y$ be a continuous map of Hausdorff spaces with dense image, show that f is an epimorphism. (Hint: Let $g, h: Y \rightarrow T$ be continuous maps of topological spaces, with T Hausdorff, show that $\{x \in Y \mid g(x) = h(x)\}$ is closed).

EXERCISE 4.10 (\star). Show that the converse holds: if $f: X \rightarrow Y$ in \mathbf{Haus} is an epimorphism, then the image of f is dense in Y .

EXERCISE 4.11 ($\star\star$). Show that the epimorphisms in \mathbf{Grp} are precisely the surjective group homomorphisms.

EXERCISE 4.12. What are the isomorphisms, monomorphisms and epimorphisms in the matrix category over a field K ? (See Example 4.10).

EXERCISE 4.13. Show that the category of fields has neither initial nor final object. Show that the category of fields of a given fixed characteristic does have an initial object.

EXERCISE 4.14 (Split monomorphisms). Let $f: X \rightarrow Y$ be a morphism in \mathcal{C} . Assume that there exists a morphism $g: Y \rightarrow X$ with $gf = \text{id}_X$. Show that f is a monomorphism. Such f is called a *split monomorphism*. Give an example of a monomorphism that is not a split monomorphism.

EXERCISE 4.15 (Split epimorphisms). Let $f: X \rightarrow Y$ be a morphism in \mathcal{C} . Assume that there exists a morphism $g: Y \rightarrow X$ with $fg = \text{id}_Y$. Show that f is an epimorphism. Such f is called a *split epimorphism*. Give an example of an epimorphism that is not a split epimorphism.

EXERCISE 4.16 (The category of G -sets). Let G be a group. A G -set is a set S together with a left action of G on S , that is, a map $G \times S \rightarrow S$, $(g, s) \mapsto gs$ satisfying $1s = s$ and $g(hs) = (gh)s$ for all $g, h \in G$ and $s \in S$. A morphism of G -sets is a map $f: S \rightarrow T$ satisfying $f(gs) = gf(s)$ for every $s \in S$ and $g \in G$. The category of G -sets is denoted ${}_G\mathbf{Set}$.

What are the final and cofinal objects of ${}_G\mathbf{Set}$? What are the monomorphisms and epimorphisms in ${}_G\mathbf{Set}$?

EXERCISE 4.17 (Homotopy category). Let \mathbf{hTop} be the *homotopy category* of topological spaces. Its objects are topological spaces, and its morphisms are *homotopy classes* of continuous maps:

$$\text{Hom}_{\mathbf{hTop}}(X, Y) := \{f: X \rightarrow Y \mid f \text{ continuous}\} / \sim$$

(Note that composition is compatible with homotopy, so that composition in \mathbf{hTop} is well-defined). Show that two topological spaces X and Y are homotopy-equivalent if and only if X and Y are isomorphic in the category \mathbf{hTop} .

EXERCISE 4.18 (\star). Let $f: X \rightarrow Y$ be a morphism in \mathcal{C} . Show that f is an isomorphism if and only if for all objects T in \mathcal{C} the map

$$\mathrm{Hom}_{\mathcal{C}}(T, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(T, Y), h \mapsto fh$$

is a bijection. (Hint, for the hard direction: use $T = Y$ to find a $g: Y \rightarrow X$ with $fg = \mathrm{id}_Y$, and use $T = X$ to deduce that $gf = \mathrm{id}_X$.)

CHAPTER 5

Functors

If in order to study groups, modules or topological spaces, one should study the homomorphisms or continuous maps between them, then to study categories one should study ‘morphisms of categories’. Such morphisms of categories are called *functors*.

1. Definition of a functor

DEFINITION 5.1. Let \mathcal{C} and \mathcal{D} be categories. A *functor* F from \mathcal{C} to \mathcal{D} , consists of the data of

- (1) for every object X in \mathcal{C} an object $F(X)$ in \mathcal{D} ,
- (2) for every morphism $f: X \rightarrow Y$ in \mathcal{C} a morphism $F(f): F(X) \rightarrow F(Y)$ in \mathcal{D}

subject to the conditions

- (F1) for every X in \mathcal{C} we have $F(\text{id}_X) = \text{id}_{F(X)}$
- (F2) for every $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ in \mathcal{C} we have $F(gf) = F(g)F(f)$.

We will write $F: \mathcal{C} \rightarrow \mathcal{D}$ to denote that F is a functor from \mathcal{C} to \mathcal{D} .

Note that if $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$ are functors, then the composite $GF: \mathcal{C} \rightarrow \mathcal{E}$ defined by $(GF)(X) := G(F(X))$ and $(GF)(f) := G(F(f))$ is also a functor.

To avoid overloading notation, we will often write FX and Ff instead of $F(X)$ and $F(f)$.

2. Many examples

EXAMPLE 5.2 (Identity). For every category \mathcal{C} there is an identity functor $\text{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$ with $\text{id}_{\mathcal{C}}(X) = X$ and $\text{id}_{\mathcal{C}}(f) = f$.

EXAMPLE 5.3 (Forgetful functors). Let R be a ring. Then we have a functor

$$F: {}_R\mathbf{Mod} \rightarrow \mathbf{Ab}$$

defined by $F(M) := M$ (as an abelian group) and $F(f) := f$. An R -module M is an abelian group equipped with extra structure, and this functor ‘forgets’ the extra structure. Other examples of forgetful functors are the obvious functors $\mathbf{Top} \rightarrow \mathbf{Set}$ (forgetting the topology), $\mathbf{Ab} \rightarrow \mathbf{Set}$ (forgetting the addition), and $\mathbf{Ring} \rightarrow \mathbf{Ab}$ (forgetting the multiplication).

EXAMPLE 5.4 (Hom functor). Let \mathcal{C} be a category in which the Hom classes are sets, and let X be an object of \mathcal{C} . Then we define a functor $F: \mathcal{C} \rightarrow \mathbf{Set}$ as follows. For an object Y in \mathcal{C} we define

$$F(Y) := \mathrm{Hom}_{\mathcal{C}}(X, Y),$$

and for a morphism $f: Y_1 \rightarrow Y_2$ in \mathcal{C} we define

$$F(f): \mathrm{Hom}_{\mathcal{C}}(X, Y_1) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, Y_2), g \mapsto fg.$$

One easily checks that F is a functor. We will denote it by $\mathrm{Hom}_{\mathcal{C}}(X, -)$.

EXAMPLE 5.5 (Abelianization of a group). If G is a group then we denote by $[G, G]$ its commutator subgroup. This is the subgroup generated by the elements $sts^{-1}t^{-1}$ with $s, t \in G$. It is a normal subgroup, and the quotient group

$$G^{\mathrm{ab}} := G/[G, G]$$

is abelian. It is called the *abelianization* of G . If $f: G \rightarrow H$ is a group homomorphism, then $f([G, G]) \subset [H, H]$, hence f induces a group homomorphism

$$f^{\mathrm{ab}}: G^{\mathrm{ab}} \rightarrow H^{\mathrm{ab}}.$$

Together, these constructions define a functor $(-)^{\mathrm{ab}}: \mathbf{Grp} \rightarrow \mathbf{Ab}$.

EXAMPLE 5.6 (Free module). Let I be a set and R be a ring. Then the set

$$R^{(I)} := \{\varphi: I \rightarrow R \mid \varphi(i) = 0 \text{ for all but finitely many } i\}$$

forms a left R -module with $(\varphi + \psi)(i) := \varphi(i) + \psi(i)$ and $(r\varphi)(i) := r\varphi(i)$. It is called the *free R -module on I* .

Note that if I is finite, then $R^{(I)}$ is free of rank the cardinality of I . If $f: I \rightarrow J$ is a map of sets, then we have an induced map

$$R^{(I)} \rightarrow R^{(J)}, \varphi \mapsto \left(j \mapsto \sum_{i \in f^{-1}(j)} \varphi(i) \right),$$

and this construction defines a functor $\mathbf{Set} \rightarrow {}_R\mathbf{Mod}$.

EXAMPLE 5.7 (Units in a ring). If $f: R \rightarrow S$ is a ring homomorphism, then f restricts to a group homomorphism $R^\times \rightarrow S^\times$, and we obtain a functor $\mathbf{Ring} \rightarrow \mathbf{Grp}$.

EXAMPLE 5.8 (The fundamental group of a pointed space). Let \mathbf{Top}_* be the category of *pointed topological spaces*. Objects in \mathbf{Top}_* are pairs (X, x) with X a topological space and $x \in X$. A morphism from (X, x) to (Y, y) is a continuous map $f: X \rightarrow Y$ such that $f(x) = y$.

Then the fundamental group defines a functor

$$\pi_1: \mathbf{Top}_* \rightarrow \mathbf{Grp}.$$

On the level of objects it is simply defined by mapping a pair (X, x) to the fundamental group $\pi_1(X, x)$. On the level of morphisms it is defined as follows. Let (X, x) and (Y, y) be pointed spaces, and let $f: X \rightarrow Y$ be a continuous map such that $f(x) = y$. Then we defined $\pi_1(f)$ by

$$\pi_1(f): \pi_1(X, x) \rightarrow \pi_1(Y, y), [\gamma] \mapsto [f \circ \gamma],$$

where $[\gamma]$ denotes the class of a loop $\gamma: [0, 1] \mapsto X$ based at x .

Note that the definition of fundamental group requires a base point. Although for a path-connected space X and points $x, y \in X$ the fundamental groups $\pi_1(X, x)$ and $\pi_1(X, y)$ are isomorphic, the isomorphism is not unique, as it depends on the choice of a path.

EXAMPLE 5.9 (Solutions to polynomial equations). Let $f_1, \dots, f_m \in \mathbf{Z}[X_1, \dots, X_n]$. Then we can consider solutions to the system of polynomial equations $f_1 = f_2 = \dots = f_m = 0$ in arbitrary commutative rings. Varying the commutative ring, one obtains a functor

$$F: \mathbf{CRing} \rightarrow \mathbf{Set}$$

defined on the level of objects by


$$R \mapsto \{x = (x_1, \dots, x_n) \in R^n \mid f_1(x) = f_2(x) = \dots = f_m(x) = 0\}.$$

and on the level of morphisms by


$$[\varphi: R \rightarrow S] \mapsto [(x_1, \dots, x_n) \mapsto (\varphi(x_1), \dots, \varphi(x_n))].$$

Indeed, if $\varphi: R \rightarrow S$ is a ring homomorphism, and if $x \in R^n$ is a solution to the system of equations, then also $(\varphi(x_1), \dots, \varphi(x_n)) \in S^n$ is a solution to the system of equations.

EXAMPLE 5.10 (Commutative diagrams as functors). Let \mathcal{C} be the small category



with $f = hg$ (and where we omitted id_x , id_y and id_z from the picture). Let \mathcal{D} be any category. Then a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ consists of objects $X := F(x)$, $Y := F(y)$, $Z := F(z)$ together with morphisms $F(f)$, $F(g)$ and $F(h)$ between them, such that $F(f) = F(h)F(g)$. In other words, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is the same as a triangular commutative diagram



in the category \mathcal{D} . With a similar construction, a commutative diagram in \mathcal{D} of any shape can be thought of as a functor from some small category \mathcal{C} to \mathcal{D} .

3. Contravariant functors

DEFINITION 5.11. Let \mathcal{C} and \mathcal{D} be categories. A *contravariant functor* from \mathcal{C} to \mathcal{D} consists of the data of

- (1) for every object X in \mathcal{C} an object $F(X)$ in \mathcal{D} ,
- (2) for every morphism $f: X \rightarrow Y$ in \mathcal{C} a morphism $F(f): F(Y) \rightarrow F(X)$ in \mathcal{D}

subject to the conditions

- (F1) for every X in \mathcal{C} we have $F(\text{id}_X) = \text{id}_{F(X)}$
- (F2') for every $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ in \mathcal{C} we have $F(gf) = F(f)F(g)$.

To stress the difference, one sometimes calls an ordinary functor a *covariant functor*.

REMARK 5.12. The only difference with the notion of a functor is that F reverses the order of composition. In other words, a contravariant functor F from \mathcal{C} to \mathcal{D} is the same as a functor $F: \mathcal{C}^{\text{opp}} \rightarrow \mathcal{D}$.

To avoid clashing notation, we will reserve the notation $F: \mathcal{C} \rightarrow \mathcal{D}$ for a functor from \mathcal{C} to \mathcal{D} , and will write $F: \mathcal{C}^{\text{opp}} \rightarrow \mathcal{D}$ for a contravariant functor from \mathcal{C} to \mathcal{D} .

EXAMPLE 5.13 (Contravariant Hom functor). Let \mathcal{C} be a category and X an object of \mathcal{C} . If $f: Y_1 \rightarrow Y_2$ is a morphism in \mathcal{C} , then we have an induced map of sets

$$\text{Hom}_{\mathcal{C}}(Y_2, X) \rightarrow \text{Hom}_{\mathcal{C}}(Y_1, X), g \mapsto gf,$$

and varying Y we obtain a contravariant functor from \mathcal{C} to **Set** given by

$$\text{Hom}_{\mathcal{C}}(-, X): \mathcal{C}^{\text{opp}} \rightarrow \mathbf{Set}, Y \mapsto \text{Hom}_{\mathcal{C}}(Y, X).$$

This is a contravariant variation on Example 5.4.

EXAMPLE 5.14. Another (related) example of a contravariant functor is the ‘dual vector space’

$$(-)^{\vee}: \mathbf{Vec}_K^{\text{opp}} \rightarrow \mathbf{Vec}_K$$

which maps a K -vector space V to its dual $V^{\vee} := \text{Hom}_K(V, K)$ and a linear map $f: V \rightarrow W$ to the induced map

$$f^{\vee}: W^{\vee} \rightarrow V^{\vee}, \varphi \mapsto \varphi \circ f.$$

EXAMPLE 5.15 (Ring of functions on a space). Yet another (related) example is the functor

$$C: \mathbf{Top}^{\text{opp}} \rightarrow \mathbf{CRing}, X \mapsto C(X) = \{\varphi: X \rightarrow \mathbf{R} \mid \varphi \text{ continuous}\},$$

mapping a topological space X to the ring of continuous \mathbf{R} -valued functions on X (with point-wise addition and multiplication). On the level of maps it is defined as follows: if $f: X \rightarrow Y$ is continuous, then we have an induced map

$$C(f): C(Y) \rightarrow C(X), \varphi \mapsto \varphi \circ f,$$

which is clearly a ring homomorphism.

4. Functors with multiple arguments

It is sometimes useful to consider functors with multiple arguments, living in various categories. This can easily be formalized using the notion of product category (see Definition 4.14). For example, a functor

$$F: \mathcal{C}_1 \times \mathcal{C}_2 \rightarrow \mathcal{D}$$

assigns to any pair of objects (X_1, X_2) with $X_i \in \text{ob } \mathcal{C}_i$ an object $F(X_1, X_2)$ in \mathcal{D} , and to any pair of morphisms (f_1, f_2) with $f_i: X_i \rightarrow Y_i$ in \mathcal{C}_i a morphism $F(f_1, f_2): F(X_1, X_2) \rightarrow F(Y_1, Y_2)$ in \mathcal{D} .

This can be combined with the notion of a contravariant functor.

EXAMPLE 5.16. A typical example is the functor

$$\text{Hom}(-, -): \mathcal{C}^{\text{opp}} \times \mathcal{C} \rightarrow \mathbf{Set},$$

which can be defined for any locally small category \mathcal{C} . If $f: X' \rightarrow X$ and $g: Y \rightarrow Y'$ are morphisms in \mathcal{C} (take note of the directions of the arrows), then we have an induced morphism

$$\text{Hom}(f, g): \text{Hom}(X, Y) \rightarrow \text{Hom}_R(X', Y'),$$

which is given by $\alpha \mapsto g\alpha f$. One says that $\text{Hom}(-, -)$ is contravariant in the first, and covariant in the second argument.

5. Morphisms of functors

DEFINITION 5.17. Let \mathcal{C} and \mathcal{D} be categories, and let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{C} \rightarrow \mathcal{D}$ be functors. A *morphism* or *natural transformation* η from F to G consists of the data of

(1) for every object X in \mathcal{C} a morphism $\eta_X: FX \rightarrow GX$ in \mathcal{D}

subject to the condition

(N1) for every morphism $f: X \rightarrow Y$ in \mathcal{C} the square

$$\begin{array}{ccc} FX & \xrightarrow{Ff} & FY \\ \downarrow \eta_X & & \downarrow \eta_Y \\ GX & \xrightarrow{Gf} & GY \end{array}$$

in \mathcal{D} commutes.

An *isomorphism* from F to G is a morphism of functors η such that η_X is an isomorphism in \mathcal{D} for every X in \mathcal{C} .

We will write $\eta: F \rightarrow G$ to denote that η is a morphism from the functor F to the functor G .

EXAMPLE 5.18 (double dual). Let K be a field and V a K -vector-space. Then we have a natural map

$$\eta_V: V \rightarrow V^{\vee\vee} = \text{Hom}_K(\text{Hom}_K(V, K), K), v \mapsto (\varphi \mapsto \varphi(v)).$$

The word ‘natural’ is usually used in an informal sense, meaning ‘not depending on the choice of a basis’. But it also has a precise mathematical meaning, namely that the collection of maps $(\eta_V)_V$ with V running over all the vector spaces forms a morphism

$$\eta: \text{id}_{\mathbf{Vec}_K} \rightarrow (-)^{\vee\vee}$$

between functors from \mathbf{Vec}_K to \mathbf{Vec}_K .

EXAMPLE 5.19. Consider the forgetful functor $F: \mathbf{Grp} \rightarrow \mathbf{Set}$. Let n be an integer. Then we have a morphism of functors $\eta: F \rightarrow F$ defined by

$$\eta_G: G \rightarrow G, g \mapsto g^n.$$

Note that $g \mapsto g^n$ is a morphism of sets, but in general not a morphism of groups.

REMARK 5.20. Note that the world of categories has three layers:

- (0) categories
- (1) functors between categories
- (2) morphisms (natural transformations) between functors

A similar picture arises in topology, where one distinguishes

- (0) topological spaces
- (1) continuous maps between spaces
- (2) homotopies between continuous maps

There is more to this than just an analogy. A fundamental construction in homotopy theory associates to a category a topological space called the *classifying space of the category*, to a functor an induced continuous map between classifying spaces, and to a natural transformation a homotopy between maps.

6. Equivalences of categories

DEFINITION 5.21. An functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is called an *equivalence* or an *equivalence of categories* if there exists a functor $G: \mathcal{D} \rightarrow \mathcal{C}$ and isomorphisms of functors

$$\epsilon: FG \xrightarrow{\sim} \text{id}_{\mathcal{D}}, \quad \eta: GF \xrightarrow{\sim} \text{id}_{\mathcal{C}}.$$

A functor G with this property is called a *quasi-inverse* of F . If there exists an equivalence from \mathcal{C} to \mathcal{D} then \mathcal{C} and \mathcal{D} are called *equivalent* categories.

REMARK 5.22. Note that this is formally very similar to the notion of a homotopy equivalence in topology. See also Remark 5.20.

Equivalent categories tend to be ‘indistinguishable’ from the point of view of category theory. See Exercises 5.17 and 5.20 for two examples. It is however often difficult to decide if a functor F is an equivalence from the definition, since it can be hard to construct a quasi-inverse functor. We end this chapter with a powerful criterion for testing if a functor F is an equivalence.

DEFINITION 5.23. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. We say that F is

(1) *full* if for every X, Y in \mathcal{C} the map

$$\mathrm{Hom}_{\mathcal{C}}(X, Y) \rightarrow \mathrm{Hom}_{\mathcal{D}}(FX, FY), f \mapsto Ff$$

is surjective;

(2) *faithful* if for every X, Y in \mathcal{C} the map

$$\mathrm{Hom}_{\mathcal{C}}(X, Y) \rightarrow \mathrm{Hom}_{\mathcal{D}}(FX, FY), f \mapsto Ff$$

is injective;

(3) *essentially surjective* if for every object Z in \mathcal{D} there is an X in \mathcal{C} such that FX and Z are isomorphic in \mathcal{D} .

THEOREM 5.24. A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is an equivalence of categories if and only if it is full, faithful and essentially surjective.

PROOF. Assume F is an equivalence of categories, with quasi-inverse G and isomorphisms

$$\epsilon: FG \rightarrow \mathrm{id}_{\mathcal{D}}, \quad \eta: GF \rightarrow \mathrm{id}_{\mathcal{C}}.$$

Then for every Y in \mathcal{D} the object $X := GY$ satisfies $FX = FGY \cong Y$ (via ϵ_Y), hence F is essentially surjective.

To see that F is faithful, let $f: X \rightarrow Y$, then we have

$$f\eta_X = \eta_Y(GFf)$$

and since η_X is an isomorphism we also have

$$f = \eta_Y(GFf)\eta_X^{-1},$$

hence clearly f is determined by Ff and $f \mapsto Ff$ is injective.

To see that F is full, let $g: FX \rightarrow FY$ be a morphism in \mathcal{D} . Then consider the morphism

$$f: X \xrightarrow{\eta_X^{-1}} GFX \xrightarrow{Gg} GFY \xrightarrow{\eta_Y} Y$$

satisfies $Ff = g$. This concludes the proof that F is full, faithful, and essentially surjective.

Conversely, assume that F is full, faithful and essentially surjective. Choose for every object X in \mathcal{D} an object GX in \mathcal{C} and an isomorphism $\alpha_Y: FGX \rightarrow X$. This is possible because of the assumption that F is essentially surjective, assuming a suitable form of the axiom of choice.

We want to make the construction $X \mapsto GX$ into a functor (which will be a quasi-inverse to F). For this, we need to define for every $f: X \rightarrow Y$ a map $Gf: GX \rightarrow GY$. Note that the map f induces a map

$$\alpha_Y^{-1} f \alpha_X: FGX \rightarrow FGY.$$

But since the functor F is full and faithful the map

$$\mathrm{Hom}_{\mathcal{C}}(GX, GY) \rightarrow \mathrm{Hom}_{\mathcal{D}}(FGX, FGY), g \mapsto Fg$$

is a bijection. Hence there exists a unique $g: GX \rightarrow GY$ with $Fg = \alpha_Y^{-1} f \alpha_X$, and we define $Gf := g$. One verifies that this G defines indeed satisfies the axioms for a functor from \mathcal{D} to \mathcal{C} .

Finally, to see that G is a quasi-inverse to F , note that the α_X define an isomorphism of functors $\alpha: FG \rightarrow \mathrm{id}_{\mathcal{D}}$. In the other direction, given an object X in \mathcal{C} we define

$$\beta_X: GFX \rightarrow X$$

to be the pre-image of α_{FX} under the bijection

$$\mathrm{Hom}_{\mathcal{C}}(GFX, X) \rightarrow \mathrm{Hom}_{\mathcal{D}}(FGFX, FX).$$

Then one verifies that the β_X define an isomorphism of functors $GF \rightarrow \mathrm{id}_{\mathcal{C}}$, and we conclude that F is an equivalence of categories. \square

EXAMPLE 5.25. Let K be a field, let \mathbf{FVec}_K be the category of finite-dimensional K -vector spaces. Let \mathcal{C} be the category of matrices over K , see Example 4.10. Consider the functor

$$F: \mathrm{Mat}_K \rightarrow \mathbf{FVec}_K$$

that maps an object n of \mathcal{C} to the vector space K^n and a matrix

$$A \in \mathrm{Hom}_{\mathcal{C}}(n, m) = \mathrm{Mat}_{m,n}(K)$$

to the corresponding linear map $K^n \rightarrow K^m$. We claim that F is an equivalence of categories.

Indeed, using Theorem 5.24 it suffices to observe that the functor is essentially surjective, since every finite-dimensional vector space is

isomorphic to K^n for some n , and that the functor is full and faithful, since for every m and n the map

$$\text{Mat}_{m,n}(K) \rightarrow \text{Hom}_K(K^n, K^m)$$

is a bijection.

Note that \mathcal{C} and \mathbf{FVec}_K are not *isomorphic* as categories. The category \mathbf{FVec}_K is much bigger than \mathcal{C} since for every n there are infinitely many n -dimensional vector spaces (all isomorphic, but not *equal*). In fact, \mathcal{C} is a small category, whereas \mathbf{FVec}_K is not.

Exercises

EXERCISE 5.1. Let \mathcal{C} and \mathcal{D} be categories, let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Let f be an isomorphism in \mathcal{C} . Show that $F(f)$ is an isomorphism in \mathcal{D} . Similarly, show that F maps split monomorphisms to split monomorphisms and split epimorphisms to split epimorphisms. (See Exercises 4.14 and 4.15).

Give examples to show that F does not necessarily map monomorphisms to monomorphisms or epimorphisms to epimorphisms.

EXERCISE 5.2. Verify the claims in Example 5.5.

EXERCISE 5.3 (Center is not a functor...). Show that there is a split monomorphism $S_2 \rightarrow S_3$ in \mathbf{Grp} . Conclude that there is no functor $F: \mathbf{Grp} \rightarrow \mathbf{Ab}$ such that for every group G we have that FG is isomorphic with the center of G .

EXERCISE 5.4 (... but it is when we restrict to isomorphisms). Let \mathbf{Grp}^\times be the category whose objects are groups, and whose morphisms are *isomorphisms* of groups (see also Exercise 4.4). Show that there is a functor $Z: \mathbf{Grp}^\times \rightarrow \mathbf{Ab}^\times$ that maps a group G to its center.

EXERCISE 5.5 (Abelianized fundamental group without base point (\star)). Let \mathcal{C} be the category of path connected topological spaces. Let P be the functor from \mathbf{Top}_\star to \mathcal{C} that maps a pair (X, x) the the path component of $x \in X$. Show that there is a functor $F: \mathcal{C} \rightarrow \mathbf{Ab}$ and an isomorphism between the functors

$$\pi_1^{\text{ab}}: \mathbf{Top}_\star \rightarrow \mathbf{Ab}, (X, x) \mapsto \pi_1(X, x)^{\text{ab}}$$

and $F \circ P$. Bonus question: show that there is *no* functor $F: \mathcal{C} \rightarrow \mathbf{Grp}$ and an isomorphism between $F \circ P$ and π_1 .

EXERCISE 5.6 (Fundamental groupoid). Let X be a topological space. Let $\Pi_1(X)$ be the category with

- (1) objects: $\text{ob } \Pi_1(X) = X$
- (2) morphisms: $\text{Hom}_{\Pi_1(X)}(x, y)$ the set of homotopy classes of paths from x to y
- (3) composition: composition of paths

Verify that this indeed defines a category. It is called the *fundamental groupoid* of X .

Assume that X is path connected, and let $x \in X$. Show that $\Pi_1(X)$ is equivalent with the category $\mathbf{B}\pi_1(X, x)$. (See Example 4.7).

EXERCISE 5.7 (Endomorphisms of identity functor on \mathbf{Ab}). Let $\eta: \text{id}_{\mathbf{Ab}} \rightarrow \text{id}_{\mathbf{Ab}}$ be a morphism of functors. Show that there is an integer n such that for every $A \in \text{ob } \mathbf{Ab}$ and for every $x \in A$ the identity $\eta_A(x) = nx$ holds.

EXERCISE 5.8. For a non-negative integer n we denote by $\text{GL}_n(R)$ the group of invertible n by n matrices with entries in R . In other words, the group of units in the ring $\text{Mat}_n(R)$, so for example $\text{GL}_1(R) = R^\times$.

Show that GL_n defines a functor from the category of commutative rings \mathbf{CRing} to the category of groups \mathbf{Grp} . Show that taking determinants defines a morphism $\det: \text{GL}_n \rightarrow \text{GL}_1$ between functors from \mathbf{CRing} to \mathbf{Grp} .

EXERCISE 5.9. Let G and H be groups, and BG and BH the corresponding one-object categories (see Example 4.7). Show that a functor $F: BG \rightarrow BH$ is the same as a group homomorphism $f: G \rightarrow H$, and a morphism of functors $\eta: F_1 \rightarrow F_2$ is the same as an element $h \in H$ such that $hf_1h^{-1} = f_2$.

EXERCISE 5.10. Let R be a ring. Recall that the *center* of a ring is the subring

$$Z(R) = \{z \in R \mid zr = rz \text{ for all } r \in R\}.$$

Denote by \mathcal{C} the category of left R -modules.

- (1) Let $z \in Z(R)$. Show that $\eta_{z,M}: M \rightarrow M, x \mapsto zx$ defines a morphism of functors $\eta_z: \text{id}_{\mathcal{C}} \rightarrow \text{id}_{\mathcal{C}}$.
- (2) Let $\eta: \text{id}_{\mathcal{C}} \rightarrow \text{id}_{\mathcal{C}}$ be a morphism of functors. Show that there is a $z \in Z(R)$ with $\eta = \eta_z$.

EXERCISE 5.11. Show that there are precisely two morphisms of functors $\text{id}_{\mathbf{Grp}} \rightarrow \text{id}_{\mathbf{Grp}}$.

EXERCISE 5.12 (Equivalence is an equivalence relation). Let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$ be equivalences of categories. Show that $GF: \mathcal{C} \rightarrow \mathcal{E}$ is an equivalence of categories.

EXERCISE 5.13. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a full and faithful functor. Let X and Y be objects in \mathcal{C} . Let $f: X \rightarrow Y$ be a morphism. Show that f is an isomorphism if and only if Ff is an isomorphism.

EXERCISE 5.14. For a category \mathcal{C} , we denote by $[\mathcal{C}]$ the class of isomorphism classes of objects in \mathcal{C} . Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Show that F induces a map $[F]: [\mathcal{C}] \rightarrow [\mathcal{D}]$. Show that if F is fully

faithful, then $[F]$ is injective, and if F is essentially surjective, then $[F]$ is surjective.

EXERCISE 5.15. Let \mathcal{C} be a locally small category in which all objects are isomorphic and in which every morphism is an isomorphism. Show that there is a group G and an equivalence of categories $\mathbf{BG} \rightarrow \mathcal{C}$.

EXERCISE 5.16. Give for every $n \in \mathbf{N}$ a category \mathcal{C}_n so that \mathcal{C}_n has exactly n objects, and all the \mathcal{C}_n are equivalent.

EXERCISE 5.17. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be an equivalence of categories. Show that \mathcal{C} has a final object if and only if \mathcal{D} has a final object.

EXERCISE 5.18. Let R and S be rings. Show that the categories ${}_R\mathbf{Mod} \times {}_S\mathbf{Mod}$ and ${}_{R \times S}\mathbf{Mod}$ are equivalent.

EXERCISE 5.19 (Morita equivalence (\star)). Let R be a ring and n a positive integer. If M is an R -module, then we can consider elements of M^n as length n column matrices with entries in M . In this way, we have an action

$$\mathrm{Mat}_n(R) \times M^n \rightarrow M^n, \left(A, \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) \mapsto A \cdot \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

This makes M^n into a left $\mathrm{Mat}_n(R)$ -module. Verify that this defines a functor

$${}_R\mathbf{Mod} \rightarrow {}_{\mathrm{Mat}_n(R)}\mathbf{Mod}, M \mapsto M^n.$$

Show that this functor is an equivalence of categories.

EXERCISE 5.20 (\star) . Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be an equivalence of categories. Let $f: X \rightarrow Y$ be a morphism in \mathcal{C} . Show that f is a monomorphism (epimorphism) if and only if $F(f)$ is a monomorphism (epimorphism).

CHAPTER 6

Tensor product

1. Tensor product of a right and a left module

DEFINITION 6.1. Let R be a ring, M a right R -module, and N a left R -module. Let A be an abelian group. A map

$$f: M \times N \rightarrow A$$

is said to be R -bilinear if for all $x, x' \in M$, $y, y' \in N$ and $r \in R$ the following hold:

- (1) $f(x + x', y) = f(x, y) + f(x', y)$
- (2) $f(x, y + y') = f(x, y) + f(x, y')$
- (3) $f(xr, y) = f(x, ry)$

Note that if $f: M \times N \rightarrow A$ is bilinear, and $h: A \rightarrow B$ is a homomorphism of abelian groups, then the composition $hf: M \times N \rightarrow B$ is also a bilinear map. The following theorem states that there is a ‘universal’ bilinear map, from which all others can be obtained by a unique composition with a homomorphism of abelian groups.

THEOREM 6.2. *Let R be a ring and let M a right R -module and N a left R -module. Then there exists an abelian group T and an R -bilinear map*

$$g: M \times N \rightarrow T$$

such that for every abelian group A and every R -bilinear map $f: M \times N \rightarrow A$ there is a unique group homomorphism $h: T \rightarrow A$ with $f = hg$:

$$\begin{array}{ccc} M \times N & \xrightarrow{g} & T \\ \downarrow f & \nearrow h & \\ A & & \end{array}$$

Moreover, the pair (T, g) is unique up to unique isomorphism in the following sense: if both (T_1, g_1) and (T_2, g_2) satisfy the above property, then there is a unique isomorphism $h: T_1 \rightarrow T_2$ such that $g_2 = hg_1$.

DEFINITION 6.3. We will call the abelian group T (unique up to unique isomorphism) the *tensor product* of M and N , and denote it by $M \otimes_R N := T$. For $x \in M$ and $y \in N$ we denote the image of (x, y) in $M \otimes_R N$ by $x \otimes y := g(x, y)$.

PROOF OF THEOREM 6.2. *Uniqueness.* This is purely formal: everything defined by a universal property is unique up to unique isomorphism, by an argument that is basically the same as the proof of Proposition 4.25: Assume (T_1, g_1) and (T_2, g_2) both satisfy the required property. Since $g_1: M \times N \rightarrow T_1$ is bilinear there is a unique map $h_1: T_2 \rightarrow T_1$ with $g_1 = h_1 g_2$. Reversing the roles of T_1 and T_2 , we get a unique map $h_2: T_1 \rightarrow T_2$. Moreover, both the compositions $h_1' h_2'$ and $h_2' h_1'$ must be the identity, so we conclude that h_1' is an isomorphism.

Existence. This part is certainly not formal! The proof is a bit messy, but in a way very natural: we just construct an abelian group T (and a map g) with all the desired properties built-in.

Let $F := \mathbf{Z}^{(M \times N)}$ be the free \mathbf{Z} -module on the set $M \times N$. Elements of F are finite integral combinations of symbols (x, y) with $x \in M$ and $y \in N$, that is, expressions of the form

$$\sum_{i=1}^n n_i (x_i, y_i)$$

with $n_i \in \mathbf{Z}$. There is a canonical map of sets

$$M \times N \rightarrow F, (x, y) \mapsto (x, y).$$

This map has no reason to be bilinear. We will force it to become bilinear by dividing out the necessary relations. Let $G \subset F$ be the subgroup generated by the elements

$$\begin{aligned} (x_1 + x_2, y) - (x_1, y) - (x_2, y), \\ (x, y_1 + y_2) - (x, y_1) - (x, y_2), \\ (xr, y) - (x, ry), \end{aligned}$$

for all $x_1, x_2, x \in M$ and $y_1, y_2, y \in N$ and $r \in R$. Let T be the quotient group F/G , and consider the composition

$$\begin{array}{ccc} M \times N & \longrightarrow & F \\ & \searrow g & \downarrow \\ & & T \end{array}$$

Then the map g is bilinear by construction.

We now show that (T, g) is a tensor product. Let $f: M \times N \rightarrow A$ be a homomorphism. Then f induces a homomorphism

$$f': F \rightarrow A, \sum n_i(x_i, y_i) \mapsto \sum n_i f(x_i, y_i).$$

Since f is bilinear, we have that f' vanishes on all the generators of G , and therefore that $f'(G) = 0$. Hence f' induces a homomorphism $h: T \rightarrow A$ with $f = hg$. To see that a map h with this property is unique, note that T is generated by the images of the elements $(x, y) \in M \times N$, and that h must send the image of (x, y) to $f(x, y)$. \square

REMARK 6.4. In practice it is often easier not to use the actual construction of the tensor product in the proof of Theorem 6.2, but only the defining universal property in the statement of Theorem 6.2, together with the fact that the tensor product is generated by the elements of the form $x \otimes y$ with $x \in M$ and $y \in N$.

REMARK 6.5. Elements of $M \otimes_R N$ are finite sums of elements of the form $x \otimes y$, but these are not independent. In fact, the map

$$M \times N \rightarrow M \otimes_R N, (x, y) \mapsto x \otimes y$$

is R -bilinear (by definition of the tensor product), so that for all $x, x_1, x_2 \in M$, $y, y_1, y_2 \in N$ and $r \in R$ the identities

$$\begin{aligned} x \otimes 0 &= 0 \\ 0 \otimes y &= 0 \\ x \otimes (y_1 + y_2) &= (x \otimes y_1) + (x \otimes y_2) \\ (x_1 + x_2) \otimes y &= (x_1 \otimes y) + (x_2 \otimes y) \\ (xr) \otimes y &= x \otimes (ry) \end{aligned}$$

hold in $M \otimes_R N$.

We end this section with a few examples of tensor products.

EXAMPLE 6.6. Let M be a right R -module, then we claim $M \otimes_R \{0\} \cong \{0\}$. Indeed, the tensor product is generated as an abelian group by the elements $x \otimes 0$, but these all are equal to 0 in $M \otimes_R \{0\}$.

EXAMPLE 6.7. We claim that for any left R -module M there is a unique isomorphism

$$f: R \otimes_R M \xrightarrow{\sim} M$$

satisfying

$$r \otimes x \mapsto rx.$$

Indeed, the map $R \times M \rightarrow M$, $(r, x) \mapsto rx$ is R -bilinear, and hence induces an R -linear homomorphism $f: R \otimes_R M \rightarrow M$ with $f(r \otimes x) = rx$. Conversely, the map $M \rightarrow R \otimes_R M$ given by $x \mapsto 1 \otimes x$ is R -linear, and is a two-sided inverse to the map f .

Of course, in the same way one can produce an isomorphism

$$M \otimes_R R \xrightarrow{\sim} M, x \otimes r \mapsto xr$$

for every *right* R -module M .

EXAMPLE 6.8. The tensor product of two non-zero modules can be zero. For example, we have

$$(\mathbf{Z}/2\mathbf{Z}) \otimes_{\mathbf{Z}} (\mathbf{Z}/3\mathbf{Z}) = 0.$$

Indeed, the tensor product is generated by elements of the form $x \otimes y$ with $x \in \mathbf{Z}/2\mathbf{Z}$ and $y \in \mathbf{Z}/3\mathbf{Z}$. But since $3x = x$ for all $x \in \mathbf{Z}/2\mathbf{Z}$ we find

$$x \otimes y = (3x) \otimes y = x \otimes (3y) = x \otimes 0 = 0.$$

Similarly we have $(\mathbf{Z}/n\mathbf{Z}) \otimes_{\mathbf{Z}} (\mathbf{Z}/m\mathbf{Z}) = 0$ whenever n and m are co-prime. See also Exercise 6.5.

2. Tensor products and bimodules

In many cases, the tensor product of two modules is not just an abelian group, but itself again a module.

DEFINITION 6.9. Let R and S be rings. An (R, S) -*bimodule* is an abelian group M equipped with operations

$$R \times M \rightarrow M, (r, x) \mapsto rx$$

and

$$M \times S \rightarrow M, (x, s) \mapsto xs$$

such that

- (B1) the first operation makes M into a left R -module
- (B2) the second operation makes M into a right S -module
- (B3) for all $r \in R$, $s \in S$ and $x \in M$ the identity $r(xs) = (rx)s$ holds in M .

A map $f: M \rightarrow N$ is called a *morphism* of (R, S) -bimodules if it is both a morphism of left R -modules and a morphism of right S -modules. We denote the category of (R, S) -bimodules by ${}_R\mathbf{Mod}_S$.

Because of axiom (B3), we can simply write rxs for $r(xs) = (rx)s$, and we will frequently describe the structure of an (R, S) -bimodule by the map

$$R \times M \times S \rightarrow M, (r, x, s) \mapsto rxs,$$

simultaneously encoding the left and right module structures.

EXAMPLE 6.10. An abelian group A has a unique structure of (\mathbf{Z}, \mathbf{Z}) -bimodule, which is given by $nam := (nm)a$ for all $n, m \in \mathbf{Z}$ and $a \in A$. It follows that a (\mathbf{Z}, \mathbf{Z}) -bimodule is the same thing as an abelian group. Similarly, an (R, \mathbf{Z}) -bimodule is the same as a left R -module, and a (\mathbf{Z}, R) -bimodule is the same as a right R -module.

EXAMPLE 6.11. R be a commutative ring and M an R -module. Then M also is an (R, R) -bimodule by setting

$$rxs := rsx$$

for all $r, s \in R$ and $x \in M$.

EXAMPLE 6.12. Let R be a ring and n a non-negative integer. Then R^n becomes an (R, R) -bimodule with

$$r(x_1, \dots, x_n)s := (rx_1s, \dots, rx_ns).$$

EXAMPLE 6.13. Let R be a commutative ring and n and m non-negative integers. Then the additive group $M = \text{Mat}_{m,n}(R)$ of n by m matrices is a $(\text{Mat}_n(R), \text{Mat}_m(R))$ -bimodule where for $A \in \text{Mat}_n(R)$, $B \in \text{Mat}_m(R)$ and $X \in M$ the element AXB of M is defined by matrix multiplication.

EXAMPLE 6.14. Let M a left S -module and N be a left R -module. Consider the group $\text{Hom}(M, N)$ of homomorphisms of abelian groups. This group carries a natural structure of (R, S) -bimodule with

$$rfs := [M \rightarrow N, x \mapsto rf(sx)]$$

for all $r \in R$, $f \in \text{Hom}(M, N)$ and $s \in S$.

PROPOSITION 6.15. Let R , S , and T be rings. Let M be an (R, S) -bimodule, and let N be an (S, T) -bimodule. Then the tensor product $M \otimes_S N$ has a unique structure of an (R, T) -bimodule satisfying

$$r(x \otimes y)t = (rx) \otimes (yt)$$

for all $r \in R$, $x \in M$, $y \in N$ and $t \in T$.

If R is a commutative ring then any R -module is canonically an (R, R) -bimodule, and we find that the tensor product of two R -modules over R is naturally an R -module. When dealing with commutative rings, there is no essential distinction between left and right modules, and we will usually treat the tensor product as an operation that produces a left R -module out of two left R -modules.

3. Tensor product as a functor

Let R be a ring and let $f: M_1 \rightarrow M_2$ be a morphism of right R -modules, and let $g: N_1 \rightarrow N_2$ be a morphism of left R -modules. Then the map

$$M_1 \times N_1 \rightarrow M_2 \otimes_R N_2, (x, y) \mapsto f(x) \otimes g(y)$$

is R -bilinear. Hence, by the universal property of the tensor product, there exists a unique homomorphism of abelian groups

$$f \otimes g: M_1 \otimes_R N_1 \rightarrow M_2 \otimes_R N_2$$

that maps an element $x \otimes y$ to $f(x) \otimes g(y)$. This upgrades the tensor product from being just a construction on pairs of modules to a functor

$$- \otimes_R -: \mathbf{Mod}_R \times {}_R\mathbf{Mod} \rightarrow \mathbf{Ab}.$$

Similarly, if R, S and T are rings, then we have a functor

$$- \otimes_S -: {}_R\mathbf{Mod}_S \times {}_S\mathbf{Mod}_T \rightarrow {}_R\mathbf{Mod}_T.$$

PROPOSITION 6.16. *Let R be a ring and let N be a left R -module. If*

$$(4) \quad M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \longrightarrow 0$$

is an exact sequence of right R -modules, then the induced sequence

$$(5) \quad M_1 \otimes_R N \xrightarrow{f \otimes \text{id}} M_2 \otimes_R N \xrightarrow{g \otimes \text{id}} M_3 \otimes_R N \longrightarrow 0$$

of abelian groups is exact. Similarly, if

$$N_1 \longrightarrow N_2 \longrightarrow N_3 \longrightarrow 0$$

is an exact sequence of left R -modules, and M a right R -module, then the induced sequence

$$M \otimes_R N_1 \longrightarrow M \otimes_R N_2 \longrightarrow M \otimes_R N_3 \longrightarrow 0$$

is exact.

The functor $- \otimes_R N$ in general does *not* preserve short exact sequences: even if

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$$

is a short exact sequence, then we only obtain a partial exact sequence

$$M_1 \otimes_R N \rightarrow M_2 \otimes_R N \rightarrow M_3 \otimes_R N \rightarrow 0.$$

See Exercise 6.13.

PROOF OF PROPOSITION 6.16. For all $x \in M_1$ and $y \in N$ we have $gf(x) \otimes y = 0$, hence $\text{im}(f \otimes \text{id}) \subset \ker(g \otimes \text{id})$. In particular, we have an induced map

$$(6) \quad \Phi: \frac{M_2 \otimes_R N}{\text{im}(f \otimes \text{id})} \longrightarrow M_3 \otimes_R N.$$

To show that the sequence (5) is exact, it suffices to show that the above map is an isomorphism. We will verify this by constructing an inverse map.

For every $x \in M_3$ choose an $x' \in M_2$ with $g(x') = x$ (note that g is surjective). We define a map

$$M_3 \otimes N \rightarrow \frac{M_2 \otimes_R N}{\text{im}(f \otimes \text{id})}, (x, y) \mapsto x' \otimes y.$$

This is well-defined, since if x'' is another element with $g(x'') = x$, then

$$(x'' \otimes y) - (x' \otimes y) = (x'' - x') \otimes y \in \text{im}(f \otimes \text{id}),$$

using the exactness of the original sequence (4). Moreover, the map is bilinear, so it induces a homomorphism

$$M_3 \otimes_R N \rightarrow \frac{M_2 \otimes_R N}{\text{im}(f \otimes \text{id})}$$

and one verifies that this is a two-sided inverse to the map Φ . \square

EXAMPLE 6.17. Proposition 6.16 can be a powerful tool in computing tensor products. As an example, let us use it to compute $M \otimes_{\mathbf{Z}} (\mathbf{Z}/n\mathbf{Z})$ for any \mathbf{Z} -module M . We have an exact sequence

$$\mathbf{Z} \xrightarrow{n} \mathbf{Z} \longrightarrow \mathbf{Z}/n\mathbf{Z} \longrightarrow 0$$

of \mathbf{Z} -modules, which induces an exact sequence

$$\mathbf{Z} \otimes_{\mathbf{Z}} M \xrightarrow{n \otimes \text{id}} \mathbf{Z} \otimes_{\mathbf{Z}} M \longrightarrow \mathbf{Z}/n\mathbf{Z} \otimes_{\mathbf{Z}} M \longrightarrow 0.$$

We have $\mathbf{Z} \otimes_{\mathbf{Z}} M \cong M$ (see Example 6.7), and the above exact sequence is isomorphic with the exact sequence

$$M \xrightarrow{n} M \longrightarrow \mathbf{Z}/n\mathbf{Z} \otimes_{\mathbf{Z}} M \longrightarrow 0,$$

from which we conclude that $\mathbf{Z}/n\mathbf{Z} \otimes_{\mathbf{Z}} M$ is isomorphic with M/nM .

4. The adjunction

Let R and S be rings. If N is an (R, S) -bimodule, and P a right S -module, then $\text{Hom}_S(N, P)$ is naturally a right R -module, with the action of R defined by:

$$rf: N \rightarrow P, x \mapsto f(rx).$$

See Exercise 6.2.

THEOREM 6.18 (Tensor-Hom adjunction). *Let R and S be rings. Let M be a right R -module, N an (R, S) -bimodule, and P be a right S -module. Then the map of abelian groups*

$$\text{Hom}_S(M \otimes_R N, P) \rightarrow \text{Hom}_R(M, \text{Hom}_S(N, P))$$

given by

$$f \mapsto (x \mapsto (y \mapsto f(x \otimes y)))$$

is an isomorphism.

PROOF. Given an R -linear map $f: M \rightarrow \text{Hom}_S(N, P)$ we obtain a map

$$M \times N \rightarrow P, (x, y) \mapsto f(x)(y)$$

which is R -bilinear, hence it defines a homomorphism

$$f': M \otimes_R N \rightarrow P$$

with the property that it maps $x \otimes y$ to $f(x)(y)$. This map is S -linear. This construction defines a homomorphism

$$\text{Hom}_R(M, \text{Hom}_S(N, P)) \rightarrow \text{Hom}_S(M \otimes_R N, P), f \mapsto f'.$$

We leave it to the reader to verify that this is a two-sided inverse to the map in the theorem. \square

5. Restriction and extension of scalars

Let $f: R \rightarrow S$ be a morphism of rings. If M is an S -module, then M also has the structure of an R -module, by $rx := f(r)x$. We obtain a functor

$${}_S\mathbf{Mod} \rightarrow {}_R\mathbf{Mod}, M \mapsto M$$

which is called *restriction of scalars*.

EXAMPLE 6.19. If $f: \mathbf{R} \rightarrow \mathbf{C}$ is the inclusion, then the corresponding restriction of scalars functor maps an n -dimensional complex vector space V , to the $2n$ -dimensional real vector space V .

EXAMPLE 6.20. If $f: \mathbf{Z} \rightarrow R$ is the canonical ring morphism from \mathbf{Z} to a commutative ring, then the restriction of scalars functor coincides with the forgetful functor

$${}_R\mathbf{Mod} \rightarrow \mathbf{Ab}, M \mapsto M$$

which maps a module to the underlying abelian group.

There is also a construction in the other direction. Let M be a (left) R -module. Note that S is an (S, R) -bimodule where $s \in S$ acts by left multiplication on S , and $r \in R$ by right multiplication with $f(r)$. Hence the tensor product $S \otimes_R M$ is a left S -module and we obtain a functor

$${}_R\mathbf{Mod} \rightarrow {}_S\mathbf{Mod}, M \mapsto S \otimes_R M$$

called *extension of scalars*.

EXAMPLE 6.21. If M is free with basis x_1, \dots, x_n , then $S \otimes_R M$ is free with basis $1 \otimes x_1, \dots, 1 \otimes x_n$. That is, every element $y \in S \otimes_R M$ can be uniquely written as

$$y = s_1 \otimes x_1 + \cdots + s_n \otimes x_n$$

with $s_i \in S$. For example, if $f: \mathbf{R} \rightarrow \mathbf{C}$ is the inclusion and $V = \mathbf{R}^n$ then we have $\mathbf{C} \otimes_{\mathbf{R}} V \cong \mathbf{C}^n$ as complex vector spaces.

PROPOSITION 6.22 (Universal property of extension of scalars). *Let $f: R \rightarrow S$ be a ring homomorphism. Let M be a left R -module and let N be a left S -module. Let $\varphi: M \rightarrow N$ be an R -linear map. Then there exists a unique S -linear map $\tilde{\varphi}: S \otimes_R M \rightarrow N$ such that the*

triangle

$$\begin{array}{ccc}
 S \otimes_R M & \overset{\tilde{\varphi}}{\dashrightarrow} & N \\
 \uparrow x \mapsto 1 \otimes x & \nearrow \varphi & \\
 M & &
 \end{array}$$

commutes.

PROOF. We first prove uniqueness. For all $x \in M$ we must have

$$\tilde{\varphi}(s \otimes x) = s\tilde{\varphi}(1 \otimes x) = s\varphi(x),$$

and since $S \otimes_R M$ is generated by elements $s \otimes x$ we see that there can be at most one $\tilde{\varphi}$. To show existence, note that

$$S \times M \rightarrow N, (s, x) \mapsto s\varphi(x)$$

is R -bilinear and hence defines a map

$$\tilde{\varphi}: S \otimes_R M \rightarrow N$$

satisfying $\tilde{\varphi}(s \otimes x) = s\varphi(x)$. For all $s, s' \in S$ and $x \in M$ we have

$$\tilde{\varphi}(s'(s \otimes x)) = \tilde{\varphi}(s's \otimes x) = s's\varphi(x) = s'(s\varphi(x)),$$

which shows that $\tilde{\varphi}$ is indeed S -linear. \square

A useful reformulation of the above proposition is the following corollary, which can also be seen as a special case of Theorem 6.18, using the canonical isomorphism $\text{Hom}_S(S, N) = N$.

COROLLARY 6.23 (Adjunction between extension and restriction of scalars). *Let $f: R \rightarrow S$ be a ring homomorphism. Let M be an R -module and let N be an S -module. Then the map*

$$\text{Hom}_S(S \otimes_R M, N) \rightarrow \text{Hom}_R(M, N), f \mapsto (x \mapsto f(1 \otimes x))$$

is a bijection. \square

Exercises

EXERCISE 6.1. Verify that the bi-module in Example 6.14 indeed satisfies the bi-module axioms.

EXERCISE 6.2. Let R and S be rings. Let M be a left R -module and N an (R, S) -bimodule. Show that $\text{Hom}_R(M, N)$ is naturally a right S -module and that $\text{Hom}_R(N, M)$ is naturally a left S -module.

EXERCISE 6.3. Let R be a ring, let M be a left R -module and let n be a non-negative integer. Show that $R^n \otimes_R M$ is isomorphic to M^n .

EXERCISE 6.4. Let R be a commutative ring. Show that $R^n \otimes_R R^m$ is isomorphic to R^{nm} as R -module.

EXERCISE 6.5. Let n and m be positive integers with greatest common divisor d . Show that

- (1) $(\mathbf{Z}/n\mathbf{Z}) \otimes_{\mathbf{Z}} (\mathbf{Z}/m\mathbf{Z}) \cong \mathbf{Z}/d\mathbf{Z}$;
- (2) $\mathbf{Q} \otimes_{\mathbf{Z}} (\mathbf{Z}/m\mathbf{Z}) \cong 0$;
- (3) $\mathbf{Q} \otimes_{\mathbf{Z}} \mathbf{Q} \cong \mathbf{Q}$.

EXERCISE 6.6. Let K be a field and $R := \text{Mat}_n(K)$ the ring of n by n matrices. Let $M = K^n$ be the right R -module of length n row vectors, and $N = K^n$ the left R -module of length n column vectors. Show that the abelian group $M \otimes_R N$ is isomorphic to K .

EXERCISE 6.7. Let R be a commutative ring and $f, g \in R$. Show that

$$R/(f) \otimes_R R/(g) \cong R/(f, g)$$

as R -modules.

EXERCISE 6.8. Let R be a ring, let $I \subset R$ be a (two-sided) ideal and let M be a left R -module. Show that $(R/I) \otimes_R M$ is isomorphic to M/IM .

EXERCISE 6.9. Let R and S be rings, let M_1 be a right R -module, M_2 an (R, S) -bimodule, and M_3 a left S -module. Show that there is a unique isomorphism

$$(M_1 \otimes_R M_2) \otimes_S M_3 \rightarrow M_1 \otimes_R (M_2 \otimes_S M_3)$$

such that

$$(x \otimes y) \otimes z \mapsto x \otimes (y \otimes z)$$

for all $x \in M_1$, $y \in M_2$ and $z \in M_3$.

EXERCISE 6.10. Let R be a ring and let M_1, M_2 be right R -modules and N a left R -module. Show that there is a unique isomorphism

$$(M_1 \oplus M_2) \otimes_R N \rightarrow (M_1 \otimes_R N) \oplus (M_2 \otimes_R N)$$

such that

$$(x, y) \otimes z \mapsto (x \otimes z, y \otimes z)$$

for all $x \in M_1, y \in M_2$ and $z \in N$.

EXERCISE 6.11. Let R be an integral domain with fraction field K . Let M be an R -module. Show that every element of $K \otimes_R M$ is of the form $\lambda \otimes x$ with $\lambda \in K$ and $x \in M$.

EXERCISE 6.12. Let R be a ring, M a right and N a left R -module. Show that there is a surjective morphism of abelian groups

$$M \otimes_{\mathbf{Z}} N \rightarrow M \otimes_R N$$

which maps $x \otimes y$ to $x \otimes y$.

EXERCISE 6.13. Consider the short exact sequence of \mathbf{Z} -modules

$$0 \rightarrow \mathbf{Z} \xrightarrow{2} \mathbf{Z} \rightarrow \mathbf{Z}/2\mathbf{Z} \rightarrow 0.$$

Show that the sequence obtained by applying the functor $-\otimes_{\mathbf{Z}} \mathbf{Z}/2\mathbf{Z}$ is *not* a short exact sequence.

EXERCISE 6.14. Let R be a ring and let $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ be a short exact sequence of right R -modules. Let N be a *free* left R -module. Show that the induced sequence

$$0 \rightarrow M_1 \otimes_R N \rightarrow M_2 \otimes_R N \rightarrow M_3 \otimes_R N \rightarrow 0$$

is a short exact sequence.

EXERCISE 6.15. Let R be a ring and let $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ be a split short exact sequence of right R -modules. Let N be a left R -module. Show that the induced sequence

$$0 \rightarrow M_1 \otimes_R N \rightarrow M_2 \otimes_R N \rightarrow M_3 \otimes_R N \rightarrow 0$$

is a short exact sequence.

EXERCISE 6.16. Let R be a principal ideal domain and let

$$M \cong R^m \oplus R/p_1^{e_1}R \oplus \cdots \oplus R/p_n^{e_n}R$$

be a finitely generated R -module (with the notation of Corollary 3.6).

(1) Let K be the fraction field of R . Compute $\dim_K(K \otimes_R M)$.

(2) Let $p \in R$ be irreducible. Compute $\dim_{R/pR}(R/pR \otimes_R M)$.

EXERCISE 6.17. Deduce the universal property of extension of scalars (Proposition 6.22) from the tensor-hom adjunction (Theorem 6.18).

EXERCISE 6.18 (Extension of scalars preserves finite generation). Let $f: R \rightarrow S$ be a morphism of rings. Let M be a finitely generated R -module. Show that $S \otimes_R M$ is a finitely generated S -module.

EXERCISE 6.19 (Restriction of scalars does not preserve finite generation). Give an example of a morphism $f: R \rightarrow S$ of commutative rings, and a finitely generated S -module M such that M is not finitely generated as an R -module.

EXERCISE 6.20. Let $f: R \rightarrow R'$ and $g: R' \rightarrow R''$ be morphisms of rings, and consider the extension of scalar functors

$${}_R\mathbf{Mod} \rightarrow {}_{R'}\mathbf{Mod} \rightarrow {}_{R''}\mathbf{Mod}.$$

Show that the composite of these functors is isomorphic to the extension of scalar functor

$${}_R\mathbf{Mod} \rightarrow {}_{R''}\mathbf{Mod}$$

for the ring morphism $gf: R \rightarrow R''$.

EXERCISE 6.21. Let $f: R \rightarrow S$ be a *surjective* morphism of rings. Show that the restriction of scalars functor

$${}_S\mathbf{Mod} \rightarrow {}_R\mathbf{Mod}, M \mapsto M$$

is full and faithful. Give an example to show that it need not be essentially surjective.

CHAPTER 7

Adjoint functors

1. Adjoint pairs of functors

DEFINITION 7.1. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{C}$ be functors between locally small categories. An *adjunction* between F and G is an isomorphism

$$\alpha: \text{Hom}_{\mathcal{D}}(F(-), -) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(-, G(-))$$

of functors $\mathcal{C}^{\text{opp}} \times \mathcal{D} \rightarrow \mathbf{Set}$ (see Example 5.16). If such an adjunction exists, we say F is *left adjoint* to G , and G is *right adjoint* to F .

In other words, an adjunction from F to G consists of the data of a bijection

$$(7) \quad \alpha_{X,Y}: \text{Hom}_{\mathcal{D}}(FX, Y) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(X, GY),$$

for every X in \mathcal{C} and Y in \mathcal{D} , such that for every $f: X_1 \rightarrow X_2$ the square

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(FX_1, Y) & \xrightarrow{\alpha_{X_1, Y}} & \text{Hom}_{\mathcal{C}}(X_1, GY) \\ \begin{array}{c} \uparrow \\ - \circ Ff \end{array} & & \begin{array}{c} \uparrow \\ - \circ f \end{array} \\ \text{Hom}_{\mathcal{D}}(FX_2, Y) & \xrightarrow{\alpha_{X_2, Y}} & \text{Hom}_{\mathcal{C}}(X_2, GY) \end{array}$$

commutes, and for every $g: Y_1 \rightarrow Y_2$ the square

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(FX, Y_1) & \xrightarrow{\alpha_{X, Y_1}} & \text{Hom}_{\mathcal{C}}(X, GY_1) \\ \begin{array}{c} \downarrow \\ g \circ - \end{array} & & \begin{array}{c} \downarrow \\ Gg \circ - \end{array} \\ \text{Hom}_{\mathcal{D}}(FX, Y_2) & \xrightarrow{\alpha_{X, Y_2}} & \text{Hom}_{\mathcal{C}}(X, GY_2) \end{array}$$

commutes. In practice, the adjunction (7) is often a canonical bijection whose functoriality is clear from the definition.

REMARK 7.2. The terminology comes from an analogy with linear algebra: if V and W are vector spaces equipped with inner products,

then linear maps $f: V \rightarrow W$ and $g: W \rightarrow V$ are called adjoint if we have

$$\langle f(v), w \rangle_W = \langle v, g(w) \rangle_V$$

for all $v \in V$ and $w \in W$.

Assume that $F: \mathcal{C} \rightarrow \mathcal{D}$ is a left adjoint of $G: \mathcal{D} \rightarrow \mathcal{C}$, with an adjunction

$$(8) \quad \alpha: \text{Hom}_{\mathcal{D}}(FX, Y) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(X, GY).$$

Taking $Y = FX$, we obtain a bijection

$$\alpha_{X, FX}: \text{Hom}_{\mathcal{D}}(FX, FX) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(X, GFX).$$

The image of id_{FX} under this map gives a map

$$\eta_X: X \rightarrow GFX$$

in \mathcal{C} , which by the functoriality of the adjunction (8) defines a morphism of functors

$$\eta: \text{id}_{\mathcal{C}} \rightarrow GF.$$

Similarly, taking $X = GY$ in (8) we obtain a morphism of functors

$$\epsilon: FG \rightarrow \text{id}_{\mathcal{D}}.$$

The morphisms of functors η and ϵ are called the *unit* and *co-unit* of the adjunction between F and G .

2. Many examples

The main reason that adjunctions between functors are interesting, is that they are ubiquitous: they arise surprisingly often in multiple branches of mathematics. Here is a short list of examples.

EXAMPLE 7.3 (Cartesian product and set of maps). Fix a set A . Then for all sets X and Y we have a canonical bijection

$$\alpha_{X, Y}: \text{Hom}(X \times A, Y) \xrightarrow{\sim} \text{Hom}(X, \text{Hom}(A, Y))$$

given by mapping a function $f: X \times A \rightarrow Y$ to the function

$$X \rightarrow \text{Hom}(A, Y), x \mapsto (a \mapsto f(x, a)).$$

An inverse is given by mapping a function $g: X \rightarrow \text{Hom}(A, Y)$ to

$$X \times A \rightarrow Y, (x, a) \mapsto g(x)(a).$$

It is easy to check that α defines an adjunction, making the functor

$$\mathbf{Set} \rightarrow \mathbf{Set}, X \mapsto X \times A$$

into a left adjoint to the functor

$$\mathbf{Set} \rightarrow \mathbf{Set}, Y \mapsto \mathrm{Hom}(A, Y).$$

The unit $\eta: \mathrm{id} \rightarrow \mathrm{Hom}(A, - \times A)$ of this adjunction is given by

$$\eta_X: X \rightarrow \mathrm{Hom}(A, X \times A), x \mapsto (a \mapsto (x, a))$$

and the co-unit $\epsilon: \mathrm{Hom}(A, -) \times A \rightarrow \mathrm{id}$ is given by

$$\epsilon_X: \mathrm{Hom}(A, X) \times A \rightarrow X, (f, a) \mapsto f(a).$$

EXAMPLE 7.4 (Tensor product and Hom). This is a variation on the previous example. Let R and S be rings, and let A be an (R, S) -bimodule. Then the functor

$$\mathbf{Mod}_R \rightarrow \mathbf{Mod}_S, M \mapsto M \otimes_R A$$

is left adjoint to the functor

$$\mathbf{Mod}_S \rightarrow \mathbf{Mod}_R, N \mapsto \mathrm{Hom}_S(A, N),$$

which comes down to the functorial isomorphism

$$\alpha_{M,N}: \mathrm{Hom}_S(M \otimes_R A, N) \xrightarrow{\sim} \mathrm{Hom}_R(M, \mathrm{Hom}_S(A, N))$$

of Theorem 6.18.

EXAMPLE 7.5 (Free module and forgetful functor). Let R be a ring. Let M be an R -module and let $R^{(I)}$ be the free R -module on a set I (see Example 5.6). Then we have a canonical map

$$\alpha_{I,M}: \mathrm{Hom}_{R\mathbf{Mod}}(R^{(I)}, M) \longrightarrow \mathrm{Hom}_{\mathbf{Set}}(I, M),$$

given by restricting a module homomorphism $\varphi: R^{(I)} \rightarrow M$ to the set of basis vectors $\{\delta_i: i \in I\}$. This map is a bijection, since a module homomorphism $R^{(I)} \rightarrow M$ is uniquely determined by the images of the basis vectors δ_i , and conversely, given a map of sets $f: I \rightarrow M$ we obtain an R -module homomorphism

$$R^{(I)} \rightarrow M, \varphi \mapsto \sum_{i \in I} \varphi(i) f(i).$$

This is just a reformulation of the familiar fact from linear algebra: to give a linear map from V to W is the same as to give the images of the vectors in a basis of V .

If we denote by

$$G: {}_R\mathbf{Mod} \rightarrow \mathbf{Set}, M \mapsto M$$

the forgetful functor (see Example 5.3) and by

$$F: \mathbf{Set} \rightarrow {}_R\mathbf{Mod}, I \mapsto R^{(I)}$$

be the free module functor, then α defines a bijection

$$\alpha_{I,M}: \mathrm{Hom}_{{}_R\mathbf{Mod}}(FI, M) \xrightarrow{\sim} \mathrm{Hom}_{\mathbf{Set}}(I, GM),$$

and one can verify directly that this defines an adjunction, making the free module functor F into a left adjoint to the forgetful functor G . The unit of this adjunction is the morphism $\eta: \mathrm{id} \rightarrow GF$ given by the function

$$\eta_I: I \rightarrow R^{(I)}, i \mapsto \delta_i,$$

for every set I .

EXAMPLE 7.6 (Discrete topology, forgetful functor, trivial topology). Any function *from* a discrete topological space is automatically continuous. Likewise, any function *to* a trivial topological space is automatically continuous. That is, we have

$$\mathrm{Hom}_{\mathbf{Top}}(X_{\mathrm{disc}}, Y) = \mathrm{Hom}_{\mathbf{Set}}(X, Y)$$

and

$$\mathrm{Hom}_{\mathbf{Set}}(X, Y) = \mathrm{Hom}_{\mathbf{Top}}(X, Y_{\mathrm{triv}}),$$

and we see that the discrete topology functor

$$\mathbf{Set} \rightarrow \mathbf{Top}, X \mapsto X_{\mathrm{disc}}$$

is left adjoint to the forgetful functor $\mathbf{Top} \rightarrow \mathbf{Set}$, and that the trivial topology functor

$$\mathbf{Set} \rightarrow \mathbf{Top}, Y \mapsto Y_{\mathrm{triv}}$$

is right adjoint to the forgetful functor.

EXAMPLE 7.7 (Frobenius reciprocity). Let k be a field, let G be a group and let $H \subset G$ be a subgroup. Then Frobenius reciprocity gives for every k -linear representation V of H and W of G a canonical isomorphism

$$\mathrm{Hom}_{k[G]}(\mathrm{Ind}_H^G V, W) \xrightarrow{\sim} \mathrm{Hom}_{k[H]}(V, \mathrm{Res}_H^G W),$$

which makes the functor Ind_H^G into a left adjoint to Res_H^G .

3. Yoneda and uniqueness of adjoints

If X is an object in \mathcal{C} , then we have a functor

$$h_X := \text{Hom}_{\mathcal{C}}(-, X): \mathcal{C}^{\text{opp}} \rightarrow \mathbf{Set}.$$

It maps an object Y to $\text{Hom}(Y, X)$, and if $f: Y_1 \rightarrow Y_2$ is a morphism in \mathcal{C} , then we obtain a morphism of sets

$$h_X(f): \text{Hom}_{\mathcal{C}}(Y_2, X) \rightarrow \text{Hom}_{\mathcal{C}}(Y_1, X), g \mapsto gf.$$

Now the functors from \mathcal{C} to \mathbf{Set} form themselves a category $\mathbf{Fun}(\mathcal{C}, \mathbf{Set})$, in which the morphisms are the natural transformations between functors. The above construction defines a functor

$$h: \mathcal{C} \rightarrow \mathbf{Fun}(\mathcal{C}^{\text{opp}}, \mathbf{Set}), X \mapsto h_X$$

On the level of morphisms it is given by sending a map $f: X \rightarrow Y$ to the natural transformation $h_f: h_X \rightarrow h_Y$ given by

$$h_{f,T}: \text{Hom}_{\mathcal{C}}(T, X) \rightarrow \text{Hom}_{\mathcal{C}}(T, Y), g \mapsto fg$$

for every T in \mathcal{C} .

THEOREM 7.8 (Yoneda's Lemma). *The functor*

$$\mathcal{C} \rightarrow \mathbf{Fun}(\mathcal{C}^{\text{opp}}, \mathbf{Set}), X \mapsto h_X$$

is fully faithful.

PROOF. In other words, we need to show that for all pairs of objects X, Y in \mathcal{C} the map

$$(9) \quad \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathbf{Fun}(\mathcal{C}^{\text{opp}}, \mathbf{Set})}(h_X, h_Y)$$

is a bijection. It is easy to define a map in the other way: Let $\varphi: h_X \rightarrow h_Y$ be a morphism of functors. Then for every T we have a map $\varphi_T: h_X(T) \rightarrow h_Y(T)$, and in particular, taking $T = X$, we find a map

$$\varphi_X: \text{Hom}_{\mathcal{C}}(X, X) \rightarrow \text{Hom}_{\mathcal{C}}(X, Y).$$

The image of id_X now gives a canonical element

$$f := \varphi_X(\text{id}_X) \in \text{Hom}_{\mathcal{C}}(X, Y)$$

a morphism $f: X \rightarrow Y$ in \mathcal{C} .

To see that this construction is a two-sided inverse to (9), note that for every $g: T \rightarrow X$ the square

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(X, X) & \xrightarrow{f \circ -} & \mathrm{Hom}_{\mathcal{C}}(X, Y) \\ \downarrow - \circ g & & \downarrow - \circ g \\ \mathrm{Hom}_{\mathcal{C}}(T, X) & \xrightarrow{f \circ -} & \mathrm{Hom}_{\mathcal{C}}(T, Y) \end{array}$$

commutes. Tracing the element id_X we find

$$\begin{array}{ccc} \mathrm{id}_X & \xrightarrow{f \circ -} & \varphi_X(\mathrm{id}_X) \\ \downarrow - \circ g & & \downarrow - \circ g \\ g & \xrightarrow{f \circ -} & \varphi_T(g) \end{array}$$

and hence that $\varphi_T(g)$ is completely determined by $f := \varphi_X(\mathrm{id}_X)$. \square

COROLLARY 7.9. *If h_X and h_Y are isomorphic functors, then X and Y are isomorphic objects in \mathcal{C} .* \square

COROLLARY 7.10 (Uniqueness of right adjoints). *If both $G_1: \mathcal{D} \rightarrow \mathcal{C}$ and $G_2: \mathcal{D} \rightarrow \mathcal{C}$ are right adjoints to a functor $F: \mathcal{C} \rightarrow \mathcal{D}$, then G_1 and G_2 are isomorphic functors.*

PROOF. Choose adjunctions between F and G_1 and between F and G_2 . Then we obtain isomorphisms

$$\mathrm{Hom}_{\mathcal{C}}(X, G_1 Y) \xleftarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(FX, Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{C}}(X, G_2 Y),$$

functorial in X and Y . Composing these, we find for all X in \mathcal{C} and Y in \mathcal{D} an isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(X, G_1 Y) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, G_2 Y).$$

Functoriality in X implies that for every Y in \mathcal{D} we find an isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(-, G_1 Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{C}}(-, G_2 Y)$$

of functors $\mathcal{C}^{\mathrm{opp}} \rightarrow \mathbf{Set}$, which by Yoneda's Lemma and Exercise 5.13 comes from a unique isomorphism

$$\gamma_Y: G_1 Y \xrightarrow{\sim} G_2 Y$$

in \mathcal{C} . Functoriality in Y implies that the collection $(\gamma_Y)_Y$ defines an isomorphism of functors

$$\gamma: G_1 \xrightarrow{\sim} G_2$$

which finishes the proof. \square

There is (of course) a dual of Yoneda's lemma. Given an object X in \mathcal{C} , consider the functor

$$h^X := \text{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathbf{Set}.$$

We have a functor

$$\mathcal{C}^{\text{opp}} \rightarrow \mathbf{Fun}(\mathcal{C}, \mathbf{Set}), X \mapsto h^X$$

On the level of morphisms it is given by sending a map $f: X \rightarrow Y$ to the natural transformation $h^f: h^Y \rightarrow h^X$ given by

$$h_T^f: \text{Hom}_{\mathcal{C}}(Y, T) \rightarrow \text{Hom}_{\mathcal{C}}(X, T), g \mapsto gf$$

for every T in \mathcal{C} .

THEOREM 7.11 (co-Yoneda's Lemma). *The functor*

$$\mathcal{C}^{\text{opp}} \rightarrow \mathbf{Fun}(\mathcal{C}, \mathbf{Set}), X \mapsto h^X$$

is fully faithful. \square

COROLLARY 7.12 (Uniqueness of left adjoints). *If both $F_1: \mathcal{C} \rightarrow \mathcal{D}$ and $F_2: \mathcal{C} \rightarrow \mathcal{D}$ are left adjoints to a functor $G: \mathcal{D} \rightarrow \mathcal{C}$, then F_1 and F_2 are isomorphic functors.* \square

Exercises

EXERCISE 7.1. Verify that the unit η and the co-unit ϵ of an adjunction are indeed morphisms of functors.

EXERCISE 7.2. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be an equivalence with quasi-inverse $G: \mathcal{D} \rightarrow \mathcal{C}$. Show that F is both left and right adjoint to G .

EXERCISE 7.3. Show that the abelianization functor $G \mapsto G^{\text{ab}}$ (see Example 5.5) is a left adjoint to the inclusion functor $\mathbf{Ab} \rightarrow \mathbf{Grp}$. What are the unit and co-unit of this adjunction?

EXERCISE 7.4. Let \mathcal{R} be the category with $\text{ob } \mathcal{R} = \mathbf{R}$ and

$$\text{Hom}_{\mathcal{R}}(x, y) = \begin{cases} \{\star\} & x \leq y \\ \emptyset & x > y \end{cases}$$

for all $x, y \in \mathbf{R}$ (see also Example 4.9). Let \mathcal{Z} be the full subcategory with $\text{ob } \mathcal{Z} = \mathbf{Z}$ and let $F: \mathcal{Z} \rightarrow \mathcal{R}$ be the inclusion functor. Does this functor have a left adjoint? And a right adjoint?

EXERCISE 7.5. Assume $F: \mathcal{C}_1 \rightarrow \mathcal{C}_2$ is left adjoint to $G: \mathcal{C}_2 \rightarrow \mathcal{C}_1$ and $F': \mathcal{C}_2 \rightarrow \mathcal{C}_3$ is left adjoint to $G': \mathcal{C}_3 \rightarrow \mathcal{C}_2$. Show that $F'F$ is left adjoint to GG' .

EXERCISE 7.6. Let $\{\star\}$ be the ‘one-point category’ consisting of a unique object \star and a unique morphism id_{\star} . Let \mathcal{C} be an arbitrary category. When does the (unique) functor $\mathcal{C} \rightarrow \{\star\}$ have a left adjoint? And a right adjoint?

EXERCISE 7.7. For a set I denote by $\mathbf{Z}[X_i \mid i \in I]$ the polynomial ring in variables (X_i) indexed by I . Elements of $\mathbf{Z}[X_i \mid i \in I]$ are finite \mathbf{Z} -linear combinations of monomials in finitely many of the variables. Verify that $I \mapsto \mathbf{Z}[X_i \mid i \in I]$ defines a functor $\mathbf{Set} \rightarrow \mathbf{CRing}$ which is left adjoint to the forgetful functor $\mathbf{CRing} \rightarrow \mathbf{Set}$.

EXERCISE 7.8. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be left adjoint to $G: \mathcal{D} \rightarrow \mathcal{C}$. Let X be a co-final object in \mathcal{C} , show that FX is cofinal in \mathcal{D} . Similarly, if Y is final in \mathcal{D} , show that GY is final in \mathcal{C} .

EXERCISE 7.9. Show that the forgetful functor $\mathbf{Top}_{\star} \rightarrow \mathbf{Top}$ has a left adjoint but not a right adjoint.

EXERCISE 7.10. Look up the definition of Stone-Čech compactification, and verify that it gives a left adjoint to the inclusion functor from the category of compact Hausdorff spaces to \mathbf{Top} .

EXERCISE 7.11. Let G be a group and let R be a ring. Show that restriction defines a bijection between the sets of

- (1) ring homomorphisms $f: \mathbf{Z}[G] \rightarrow R$
- (2) group homomorphisms $G \rightarrow R^\times$.

Interpret this bijection as an adjunction between functors between the categories of groups and rings.

EXERCISE 7.12 (Triangle identities (\star)). Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a left adjoint to $G: \mathcal{D} \rightarrow \mathcal{C}$, with unit $\eta: \text{id}_{\mathcal{C}} \rightarrow GF$ and co-unit $\epsilon: FG \rightarrow \text{id}_{\mathcal{D}}$. Show that the diagrams

$$\begin{array}{ccc} FX & \xrightarrow{F\eta_X} & FGFX \\ & \searrow \text{id} & \downarrow \epsilon_{FX} \\ & & FX \end{array} \qquad \begin{array}{ccc} GY & \xrightarrow{\eta_{GY}} & GFGY \\ & \searrow \text{id} & \downarrow G\epsilon_Y \\ & & GY \end{array}$$

commute for every X in \mathcal{C} and Y in \mathcal{D} .

EXERCISE 7.13. A functor $F: \mathcal{C} \rightarrow \mathbf{Set}$ is called *co-representable* if there exists an object X in \mathcal{C} with $h^X \cong F$. We say that F is co-represented by X . Let $f_1, \dots, f_m \in \mathbf{Z}[X_1, \dots, X_n]$. Show that the functor

$$\mathbf{CRing} \rightarrow \mathbf{Set}, R \mapsto \{x \in R^n \mid f_1(x) = \dots = f_m(x) = 0\}$$

of Example 5.9 is co-representable.

EXERCISE 7.14 (\star) . Show that the functor

$$\mathbf{GL}_n: \mathbf{CRing} \rightarrow \mathbf{Set}, R \mapsto \mathbf{GL}_n(R)$$

of Exercise 5.8 is co-representable. (Hint: first show that the functor $\mathbf{GL}_1: R \mapsto R^\times$ is isomorphic to h^{R_1} with $R_1 = \mathbf{Z}[X, Y]/(XY - 1)$.) Let R_n be the commutative ring such that $\mathbf{GL}_n \cong h^{R_n}$. By the co-Yoneda lemma there is a unique ring homomorphism $R_1 \rightarrow R_n$ inducing the natural transformation $\det: \mathbf{GL}_n \rightarrow \mathbf{GL}_1$. Describe this ring homomorphism explicitly.

EXERCISE 7.15 (\star) . For topological spaces A and Y define $C(A, Y)$ to be the set of continuous maps from A to Y . For every compact $K \subset A$ and open $U \subset Y$ let $C_{K,U} \subset C(A, Y)$ be the subset consisting of those $f: A \rightarrow Y$ with $f(K) \subset U$. We give $C(A, Y)$ the topology generated by the subsets $C_{K,U}$. (It is known as the compact-open topology).

Show that if A is compact and Hausdorff, then the functor

$$\mathbf{Top} \rightarrow \mathbf{Top}: X \mapsto X \times A$$

(giving $X \times A$ the product topology) is left adjoint to the functor

$$\mathbf{Top} \rightarrow \mathbf{Top}: Y \mapsto C(A, Y).$$

CHAPTER 8

Limits and colimits

1. Product and coproduct

DEFINITION 8.1. Let \mathcal{C} be a category, let X and Y be objects in \mathcal{C} . A *product* of X and Y consists of the data of

- (1) an object P
- (2) morphisms $\pi_X: P \rightarrow X$ and $\pi_Y: P \rightarrow Y$

such that for all objects T in \mathcal{C} , and for all $f: T \rightarrow X$ and $g: T \rightarrow Y$ there is a unique map $h: T \rightarrow P$ making the following diagram commute

$$\begin{array}{ccc} & & X \\ & \nearrow f & \nearrow \\ T & \overset{h}{\dashrightarrow} P & \nearrow \pi_X \\ & \searrow g & \searrow \pi_Y \\ & & Y \end{array}$$

A product need not exist, but if it does, it is unique up to unique isomorphism. In particular, we will refer to any product of X and Y as *the product* of X and Y . We will usually denote it by $X \times Y$, omitting the morphisms π_X and π_Y . The proof of uniqueness is formal, and essentially identical to the proofs of Proposition 4.25 and the uniqueness part of Theorem 6.2. We repeat the argument for the last time.

PROPOSITION 8.2. Let (P, π_X, π_Y) and (P', π'_X, π'_Y) be products of X and Y . Then there exists a unique isomorphism $h: P \rightarrow P'$ such

that the diagram

$$\begin{array}{ccc}
 & X & \\
 \pi_X \nearrow & & \nwarrow \pi'_X \\
 P & \overset{h}{\dashrightarrow} & P' \\
 \pi_Y \searrow & & \swarrow \pi'_Y \\
 & Y &
 \end{array}$$

commutes.

PROOF OF PROPOSITION 8.2. Since P' is a product, there exists a unique $h: P \rightarrow P'$ making the diagram commute. Likewise, since P is a product, there exists a unique $h': P' \rightarrow P$ making the diagram commute. By the same reasoning, there are unique maps i and i' making the diagrams

$$\begin{array}{ccc}
 & X & \\
 \pi_X \nearrow & & \nwarrow \pi_X \\
 P & \overset{i}{\dashrightarrow} & P \\
 \pi_Y \searrow & & \swarrow \pi_Y \\
 & Y &
 \end{array}
 \qquad
 \begin{array}{ccc}
 & X & \\
 \pi'_X \nearrow & & \nwarrow \pi'_X \\
 P' & \overset{i'}{\dashrightarrow} & P' \\
 \pi'_Y \searrow & & \swarrow \pi'_Y \\
 & Y &
 \end{array}$$

commute. Combining both parts, we see that both $i = \text{id}_P$ and $i = h'h$ make the first diagram commute, and that both $i = \text{id}_{P'}$ and $i = hh'$ make the second diagram commute. By unicity we have $h'h = \text{id}_P$ and $hh' = \text{id}_{P'}$, hence h is an isomorphism. \square

EXAMPLES 8.3. In **Set** the product is the cartesian product, together with the usual projections. In **Top** the product is the cartesian product equipped the product topology, together with the usual projections. The product of two rings R and S in **Ring** is the product ring $R \times S$. The product of two modules in $R\mathbf{Mod}$ is the cartesian product $M \times N$.

The dual notion of a product is a coproduct (sometimes called sum).

DEFINITION 8.4. Let \mathcal{C} be a category, let X and Y be objects in \mathcal{C} . A *coproduct* of X and Y consists of the data of

- (1) an object S
- (2) morphisms $\iota_X: X \rightarrow S$ and $\iota_Y: Y \rightarrow S$

such that for all objects T in \mathcal{C} , and for all $f: X \rightarrow T$ and $g: Y \rightarrow T$ there is a unique map $h: S \rightarrow T$ making the diagram

$$\begin{array}{ccc}
 X & & T \\
 \searrow & \curvearrowright f & \\
 & S \dashrightarrow & \\
 \swarrow & \curvearrowleft g & \\
 Y & &
 \end{array}$$

commute.

By the same argument as before, a coproduct, if it exists, is unique up to isomorphism. We talk about *the* coproduct, and denote it with $X \amalg Y$.

EXAMPLE 8.5. Let X and Y be sets. Then the coproduct of X and Y in **Set** is the disjoint union $X \amalg Y$, together with the canonical inclusions $\iota_X: X \rightarrow X \amalg Y$ and $\iota_Y: Y \rightarrow X \amalg Y$.

Similarly, the coproduct in **Top** of topological spaces X and Y is the disjoint union $X \amalg Y$, with the natural topology (in which X and Y are both open and closed).

EXAMPLE 8.6. Let R be a ring. Then the coproduct of R -modules M and N is the direct sum $M \oplus N$, and hence in this case the product and the coproduct coincide (although the former is considered together with the projections, and the latter with the inclusions).

More generally one can define products and coproducts of arbitrary (possibly infinite) families of objects.

DEFINITION 8.7. The *product* of a family $(X_i)_{i \in I}$ of objects in \mathcal{C} is an object $\prod_{i \in I} X_i$ together with maps $\pi_n: \prod_{i \in I} X_i \rightarrow X_n$ such that for every T and for every collection of morphisms $f_i: T \rightarrow X_i$ there is a unique $h: T \rightarrow \prod_{i \in I} X_i$ such that $f_i = \pi_i h$ for all $i \in I$.

DEFINITION 8.8. The *coproduct* of a family $(X_i)_{i \in I}$ of objects in \mathcal{C} is an object $\amalg_{i \in I} X_i$ together with maps $\iota_n: X_n \rightarrow \amalg_{i \in I} X_i$ such that for every T and for every collection of morphisms $f_i: X_i \rightarrow T$ there is a unique $h: \amalg_{i \in I} X_i \rightarrow T$ such that $f_i = h \iota_i$ for all $i \in I$.

Again, products and coproducts of arbitrary families need not exist, but if they do they are unique up to unique isomorphism.

EXAMPLE 8.9 (Product and coproduct of modules). Let R be a ring and $(M_i)_{i \in I}$ a collection of R -modules. Then the product of (M_i) is the R -module

$$\prod_{i \in I} M_i = \{(x_i)_{i \in I} \mid x_i \in M_i\}$$

together with the projection maps $\pi_n: \prod_{i \in I} M_i \rightarrow M_n$. The universal property of products gives a natural (*i.e.* functorial) bijection

$$\mathrm{Hom}_R(N, \prod_{i \in I} M_i) \xrightarrow{\sim} \prod_{i \in I} \mathrm{Hom}_R(N, M_i).$$

The coproduct of $(M_i)_i$ is the direct sum $\bigoplus_{i \in I} M_i$ together with the inclusion maps $\iota_n: M_n \rightarrow \bigoplus_{i \in I} M_i$ given by

$$\iota_n(x)_i = \begin{cases} x & i = n \\ 0 & i \neq n \end{cases}$$

for all $x \in M_n$ and $i \in I$. Indeed, the natural bijection

$$\mathrm{Hom}_R(\bigoplus_{i \in I} M_i, N) \xrightarrow{\sim} \prod_{i \in I} \mathrm{Hom}_R(M_i, N)$$

of Exercise 1.17 shows that for every collection of morphisms $(f_i: M_i \rightarrow N)_i$ there is a unique morphism $h: \bigoplus_{i \in I} M_i \rightarrow N$ with $h\iota_i = f_i$.

2. Pushout and pullback

DEFINITION 8.10. Let \mathcal{C} be a category and let $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ be morphisms in \mathcal{C} . The *pull-back* or *fibred product* of f and g consists of

- (1) an object P in \mathcal{C}
- (2) morphisms $\pi_X: P \rightarrow X$ and $\pi_Y: P \rightarrow Y$

such that $f\pi_X = g\pi_Y$ as maps $P \rightarrow Z$, and such that for every object T and for every pair of maps $s: T \rightarrow X$, $t: T \rightarrow Y$ with $fs = gt$ there is a unique morphism $h: T \rightarrow P$ making the diagram

$$\begin{array}{ccccc}
 & & & X & \\
 & & s \nearrow & \nearrow & \\
 & & & \pi_X \nearrow & \\
 T & \xrightarrow{h} & P & & Z \\
 & & \searrow \pi_Y & \searrow & \\
 & & & Y & \\
 & & t \searrow & \searrow & \\
 & & & &
 \end{array}$$

commute.

If it exists, the pull-back is unique up to unique isomorphism. It is usually denoted by $P = X \times_Z Y$, but care should be taken since the pull-back does depend on the maps f and g . Alternatively, one says that the square

$$\begin{array}{ccc} P & \xrightarrow{\pi_Y} & Y \\ \pi_X \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

is *cartesian* if (P, π_X, π_Y) is the fibered product of f and g .

EXAMPLE 8.11. In the category of sets, the fiber product of any pair of maps $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ exists. It is given by

$$X \times_Z Y = \{(x, y) \in X \times Y \mid f(x) = g(y)\},$$

together with the projection maps. Indeed, if $s: T \rightarrow X$ and $t: T \rightarrow Y$ satisfy $fs = gt$, then the map

$$h: T \rightarrow X \times_Z Y, u \mapsto (s(u), t(u)).$$

is the unique map making the diagram of the definition commute.

In the case that $X = \{\star\}$ and f the map $\star \mapsto z$, then we find

$$\{\star\} \times_Z Y = g^{-1}(z),$$

the fiber of Y over z .

In the case that X, Y are subsets of Z and f and g are the respective inclusions we find $X \times_Z Y = X \cap Y$.

EXAMPLE 8.12. Similarly, in **Top** we have

$$X \times_Z Y = \{(x, y) \in X \times Y \mid f(x) = g(y)\}$$

with the induced topology from the product topology on $X \times Y$.

EXAMPLE 8.13. Similarly, if $f_1: M_1 \rightarrow N$ and $f_2: M_2 \rightarrow N$ are R -module homomorphisms, then

$$\{(x_1, x_2) \in M_1 \times M_2 \mid f_1(x_1) = f_2(x_2)\}$$


is a sub- R -module of $M_1 \times M_2$, and one verifies that it is the pull-back of f_1 and f_2 .

The dual notion of pullback/fibered product is pushout/fibered co-product. The definition is obtained by reversing all the arrows.

DEFINITION 8.14. Let \mathcal{C} be a category and let $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ be morphisms in \mathcal{C} . The *pushout* or *fiber coproduct* or *fiber sum* of f and g consists of

- (1) an object S in \mathcal{C}
- (2) morphisms $\iota_X: X \rightarrow S$ and $\iota_Y: Y \rightarrow S$


such that $\iota_X f = \iota_Y g$ as maps $Z \rightarrow S$, and such that for every object T and for every pair of maps $s: X \rightarrow T$, $t: Y \rightarrow T$ with $sf = tg$ there is a unique morphism $h: S \rightarrow T$ making the diagram



commute.

If it exists, the pushout is unique up to unique isomorphism.

EXAMPLE 8.15. Pushouts exist in **Set**, and can be constructed as follows. Let $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ be functions. Consider the equivalence relation \sim on the disjoint union $X \amalg Y$ generated by $f(z) \sim g(z)$. Let S be the quotient $(X \amalg Y)/\sim$. Then we claim that S is the pushout of f and g . Indeed, assume that we have a commutative diagram



Then there is only one possibility for the map h :

$$h: S \rightarrow T, \begin{cases} [x] \mapsto s(x) & x \in X \\ [y] \mapsto t(y) & y \in Y \end{cases}$$

This is indeed well-defined, since for any $z \in Z$ we have $s(f(z)) = t(g(z))$.

EXAMPLE 8.16 (Glueing). The same construction as in **Set** defines a pushout in **Top**, it suffices to put the quotient topology on $X \amalg Y / \sim$. This construction is particularly useful in topology when both $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ are injective. In this case it constructs a space S by ‘gluing’ X and Y along their common subspace Z .

Conversely, if S is a topological space and X and Y subspaces with $S = X \cup Y$, then S is the pushout of the inclusion maps $X \cap Y \rightarrow X$ and $X \cap Y \rightarrow Y$.

3. Equalizer and co-equalizer

DEFINITION 8.17. Let \mathcal{C} be a category, let X and Y be objects in \mathcal{C} and let $f: X \rightarrow Y$ and $g: X \rightarrow Y$ be morphisms in \mathcal{C} . The *equalizer* of f and g is a pair (E, ι) consisting of

- (1) an object E in \mathcal{C} ,
- (2) a morphism $\iota: E \rightarrow X$ in \mathcal{C} ,

such that $f\iota = g\iota$, and such that for every morphism $t: T \rightarrow X$ with $ft = gt$ there is a unique $h: T \rightarrow E$ making the diagram

$$\begin{array}{ccccc} T & & & & \\ \downarrow h & \searrow t & & & \\ E & \xrightarrow{\iota} & X & \xrightleftharpoons[g]{f} & Y \end{array}$$

commute.

The equalizer of f and g , if it exists, is unique up to unique isomorphism.

EXAMPLE 8.18. In **Set** the equalizer of $f, g: X \rightarrow Y$ is given by $E = \{x \in X \mid f(x) = g(x)\}$ and $\iota: E \rightarrow X$ the inclusion. Also in **Ring**, **CRing**, **Top**, **Grp**, ${}_R\mathbf{Mod}$ equalizers exist, and are given by the same recipe. For example, the equalizer of a pair of ring homomorphisms $f, g: R \rightarrow S$ is the subring

$$\{r \in R \mid f(r) = g(r)\} \subset R,$$

together with its inclusion in R .

EXAMPLE 8.19. In ${}_R\mathbf{Mod}$ the equalizer of $f, g: M \rightarrow N$ is nothing but $\ker(f - g)$. In particular, taking $g = 0$ we see that a kernel of an R -module homomorphism is a special case of an equalizer.

The dual notion of equalizer is co-equalizer.

DEFINITION 8.20. Let \mathcal{C} be a category, let X and Y be objects in \mathcal{C} and let $f: X \rightarrow Y$ and $g: X \rightarrow Y$ be morphisms in \mathcal{C} . The *co-equalizer* of f and g is a pair (C, π) consisting of

- (1) an object C in \mathcal{C} ,
- (2) a morphism $\pi: Y \rightarrow C$ in \mathcal{C} ,

such that $\pi f = \pi g$, and such that for every morphism $t: Y \rightarrow T$ with $tf = tg$ there is a unique $h: C \rightarrow T$ making the diagram

$$\begin{array}{ccc} X & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & Y & \xrightarrow{\pi} & C \\ & & & \searrow t & \downarrow h \\ & & & & T \end{array}$$

commute.

The co-equalizer of f and g , if it exists, is unique up to unique isomorphism.

EXAMPLE 8.21. In **Set**, the co-equalizer of $f, g: X \rightarrow Y$ is the quotient $E = Y/\sim$ of Y by the equivalence relation generated by $f(x) \sim g(x)$ for all $x \in X$.

EXAMPLE 8.22. In $R\mathbf{Mod}$ the co-equalizer of $f, g: M \rightarrow N$ is $\text{coker}(f - g)$. In particular, taking $g = 0$ we see that cokernels of R -linear maps are examples of co-equalizers.

4. Limits and colimits

The above six constructions are examples of two dual general classes of categorical constructions called limits and colimits.

Let \mathcal{I} be a small category (see Definition 4.11), let \mathcal{C} be a category, and let $X: \mathcal{I} \rightarrow \mathcal{C}$ be a functor. We will often denote the image of an object $i \in \mathcal{I}$ by X_i .

It is useful to think of the functor $X: \mathcal{I} \rightarrow \mathcal{C}$ informally as a diagram in \mathcal{C} , indexed by \mathcal{I} . For example, if \mathcal{I} is the three-object category

$$\begin{array}{ccc} 1 & & \\ & \searrow \varphi & \\ & & 2 \\ & \nearrow \psi & \\ 3 & & \end{array}$$

then a functor $X: \mathcal{I} \rightarrow \mathcal{C}$ can be thought of as a diagram

$$\begin{array}{ccc} X_1 & & \\ & \searrow^{X(\varphi)} & \\ & & X_2 \\ & \nearrow_{X(\psi)} & \\ X_3 & & \end{array}$$

of objects and morphisms in \mathcal{C} . See also Example 5.10.

DEFINITION 8.23. The *limit* of a functor $X: \mathcal{I} \rightarrow \mathcal{C}$ consists of

- (1) an object $\lim_{\mathcal{I}} X$ in \mathcal{C}
- (2) for every object i in \mathcal{I} a morphism $\pi_i: \lim_{\mathcal{I}} X \rightarrow X_i$

such that

- (1) for every $\varphi: i \rightarrow j$ in \mathcal{I} we have $\pi_j = X(\varphi) \circ \pi_i$
- (2) for every T in \mathcal{C} and for every collection of morphisms $t_i: T \rightarrow X_i$ satisfying $t_j = X(\varphi) \circ t_i$ for all $\varphi: i \rightarrow j$, there is a unique $h: T \rightarrow \lim_{\mathcal{I}} X$ with $t_i = \pi_i h$ for all i .

Of course, the limit of $\mathcal{I} \rightarrow \mathcal{C}$, if it exists, is unique up to unique isomorphism. The limit is sometimes written $\lim_i X_i$ or $\lim_{\mathcal{I}} X_i$, but one should be careful to remember that it depends on the full functor X , and not just on the objects $(X_i)_{i \in \text{ob } \mathcal{I}}$.

EXAMPLES 8.24. Taking for \mathcal{I} the three categories

$$\begin{array}{ccc} & & 1 \\ & & \searrow \\ 1 & & 2, \\ & & \nearrow \\ & & 3 \end{array} \qquad 1 \rightrightarrows 2$$

recovers the notions of product, fibered product and equalizer. If we take for \mathcal{I} the empty category, then the limit (of the unique functor $\emptyset \rightarrow \mathcal{C}$) is the final object of \mathcal{C} (if it exists).

There is no surprise in the definition of colimit:

DEFINITION 8.25. The *colimit* of a functor $X: \mathcal{I} \rightarrow \mathcal{C}$ consists of

- (1) an object $\text{colim}_{\mathcal{I}} X$ in \mathcal{C} ,
- (2) for every object i in \mathcal{I} a morphism $\iota_i: X_i \rightarrow \text{colim}_{\mathcal{I}} X$

such that

- (1) for every $\varphi: i \rightarrow j$ in \mathcal{I} we have $\iota_j \circ X(\varphi) = \iota_i$,
 (2) for every T in \mathcal{C} and for every collection of morphisms $t_i: X_i \rightarrow T$ satisfying $t_j \circ X(\varphi) = t_i$ for all $\varphi: i \rightarrow j$, there is a unique $h: \operatorname{colim}_{\mathcal{I}} X \rightarrow T$ with $t_i = h\iota_i$ for all i .

If it exists, it is unique up to unique isomorphism.

EXAMPLES 8.26. Taking for \mathcal{I} the opposites of the three categories in 8.24 one recovers the notions of coproduct, fibered coproduct and co-equalizer. If we take for \mathcal{I} the empty category, then the colimit (of the unique functor $\emptyset \rightarrow \mathcal{C}$) is the cofinal object of \mathcal{C} (if it exists).

We end with an example of an infinite diagram in **Set** and its limit and colimit.

EXAMPLE 8.27. Let \mathcal{I} be the category with $\operatorname{ob}\mathcal{I} = \mathbf{N}$ and such that for all $i < j$ we have $\operatorname{Hom}(i, j) = \emptyset$, and for all $i \geq j$ we have $\operatorname{Hom}(i, j) = \{\star\}$. In a picture:

$$0 \longleftarrow 1 \longleftarrow 2 \longleftarrow 3 \longleftarrow \cdots .$$

Let $S_0 \supset S_1 \supset S_2 \supset \cdots$ be a decreasing chain of sets. Then this defines a functor

$$S: \mathcal{I} \rightarrow \mathbf{Set}, i \mapsto S_i$$

which for $i \geq j$ maps the unique map $i \rightarrow j$ to the inclusion $S_i \hookrightarrow S_j$.

To give a collection of maps $t_i: T \rightarrow S_i$ such that for all $i \geq j$ the diagram

$$\begin{array}{ccc} T & & \\ \downarrow t_j & \searrow t_i & \\ S_j & \longleftarrow & S_i \end{array}$$

commutes, is the same as to give a map $t: T \rightarrow \bigcap_i S_i$. From this it follows easily that $\lim_i S_i = \bigcap_i S_i$.

For the colimit, note that a collection of maps $t_i: S_i \rightarrow T$ such that for all $i \geq j$ the diagram

$$\begin{array}{ccc} S_j & \longleftarrow & S_i \\ & \searrow t_j & \downarrow t_i \\ & & T \end{array}$$

commutes is completely determined by $t_0: S_0 \rightarrow T$ (since t_i must be the restriction of t_0 to the subset $S_i \subset S_0$), and one easily verifies that $\operatorname{colim}_i S_i = S_0$.

5. Yoneda and limits and colimits of sets

The category **Set** has all limits and colimits.

PROPOSITION 8.28. *Let \mathcal{I} be a small category and let $X: \mathcal{I} \rightarrow \mathbf{Set}$, $i \mapsto X_i$ be a functor. Then $\lim_{\mathcal{I}} X_i$ exists, and is given by*

$$\{(x_i)_i \in \prod_{i \in \operatorname{ob} \mathcal{I}} X_i \mid X(\varphi)(x_i) = x_j \text{ for all } \varphi: i \rightarrow j\}$$

together with the projection maps to the sets X_i .

PROOF. The proof is a straightforward verification that the set $L := \{(x_i)_i \in \prod_i X_i \mid \dots\}$ described in the proposition, with the projection maps

$$\pi_n: L \rightarrow X_n, (x_i)_i \mapsto x_n$$

satisfies the definition of a limit. The first property holds by construction: for every $x \in L$ and $\varphi: i \rightarrow j$ in \mathcal{I} we have

$$\pi_j(x) = x_j = X_\varphi(x_i) = (X_\varphi \circ \pi_i)(x),$$

hence $\pi_j = X_\varphi \circ \pi_i$. For the second property, assume that $(T, (t_i: T \rightarrow X_i)_i)$ satisfies $t_j = X_\varphi \circ t_i$ for every $\varphi: i \rightarrow j$. Then the map

$$h: T \rightarrow L, x \mapsto (t_i(x))_i$$

is well-defined (that is, $h(x)$ lands in $L \subset \prod_i X_i$), and clearly is the unique map such that $t_i = \pi_i h$ for every i . \square

PROPOSITION 8.29. *Let \mathcal{I} be a small category and let $X: \mathcal{I} \rightarrow \mathbf{Set}$, $i \mapsto X_i$ be a functor. Consider on the disjoint union $\coprod_{i \in \operatorname{ob} \mathcal{I}} X_i$ the equivalence relation \sim generated by $x_i \sim X(\varphi)(x_i)$ for all $\varphi: i \rightarrow j$ and all $x_i \in X_i$. Then $\operatorname{colim}_{\mathcal{I}} X_i$ exists and is given by*

$$\operatorname{colim}_{\mathcal{I}} X = \left(\prod_{i \in \operatorname{ob} \mathcal{I}} X_i \right) / \sim$$

together with the compositions

$$\iota_i: X_i \hookrightarrow \prod_{i \in \operatorname{ob} \mathcal{I}} X_i \twoheadrightarrow \operatorname{colim}_{\mathcal{I}} X.$$

PROOF. This is shown by a direct verification, somewhat similar to the proof of Proposition 8.28. See also Exercise 8.19. \square

Using the explicit descriptions of limits of sets, we can now rephrase the universal property for limits and colimits in an arbitrary category:

THEOREM 8.30. *Let \mathcal{I} be a small category and let $X: \mathcal{I} \rightarrow \mathcal{C}$ be a functor. Let L be an object of \mathcal{C} . Then $\lim_{\mathcal{I}} X$ exists and is isomorphic to L if and only if there is an isomorphism*

$$\mathrm{Hom}_{\mathcal{C}}(-, L) \xrightarrow{\sim} \lim_{\mathcal{I}} \mathrm{Hom}_{\mathcal{C}}(-, X_i),$$

of functors $\mathcal{C}^{\mathrm{opp}} \rightarrow \mathbf{Set}$.

Note that by Yoneda (Theorem 7.8) the above theorem completely characterizes $\lim_{\mathcal{I}} X$.

PROOF OF THEOREM 8.30. Let T be an object in \mathcal{C} . Then by the universal property of the limit in \mathcal{C} we have a natural bijection

$$\mathrm{Hom}_{\mathcal{C}}(T, \lim_i X_i) \xrightarrow{\sim} \{(t_i: T \rightarrow X_i)_i \mid \forall \varphi: i \rightarrow j, X(\varphi) \circ t_i = t_j\},$$

and by the explicit description of limit of sets in Proposition 8.28 the latter equals $\lim_i \mathrm{Hom}_{\mathcal{C}}(T, X_i)$. The bijection is functorial in T , and hence defines an isomorphism of functors. Conversely, if there is a functorial bijection

$$\mathrm{Hom}_{\mathcal{C}}(T, L) \xrightarrow{\sim} \{(t_i: T \rightarrow X_i)_i \mid \forall \varphi: i \rightarrow j, X(\varphi) \circ t_i = t_j\},$$

then L satisfies the universal property of the limit. \square

THEOREM 8.31. *Let \mathcal{I} be a small category and let $X: \mathcal{I} \rightarrow \mathcal{C}$ be a functor. Let C be an object of \mathcal{C} . Then $\mathrm{colim}_{\mathcal{I}} X$ exists and is isomorphic to C if and only if there is an isomorphism*

$$\mathrm{Hom}_{\mathcal{C}}(C, -) = \lim_{\mathcal{I}^{\mathrm{opp}}} \mathrm{Hom}_{\mathcal{C}}(X_i, -)$$

of functors $\mathcal{C} \rightarrow \mathbf{Set}$. \square

Note that by co-Yoneda (Theorem 7.11) the above theorem completely characterizes $\mathrm{colim}_{\mathcal{I}} X$.

6. Adjoint functors and limits

THEOREM 8.32 (Left adjoints commute with colimits). *Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a left adjoint to $G: \mathcal{D} \rightarrow \mathcal{C}$. Let $X: \mathcal{I} \rightarrow \mathcal{C}$ be a functor, and suppose that $\operatorname{colim}_{\mathcal{I}} X$ exists in \mathcal{D} . Then $\operatorname{colim}_{\mathcal{I}}(FX)$ exists and*

$$\operatorname{colim}_{\mathcal{I}}(FX) \cong F(\operatorname{colim}_{\mathcal{I}} X)$$

in \mathcal{D} .

Informally, we say that ‘ F commutes with colimits’.

PROOF OF THEOREM 8.32. Using Theorem 8.31 and the definition of adjoint functors we find for every T in \mathcal{D} a chain of isomorphisms

$$\begin{aligned} \operatorname{Hom}_{\mathcal{D}}(F(\operatorname{colim}_{\mathcal{I}} X_i), T) &\cong \operatorname{Hom}_{\mathcal{C}}(\operatorname{colim}_{\mathcal{I}} X_i, GT) \\ &\cong \lim_{\mathcal{I}^{\text{opp}}} \operatorname{Hom}_{\mathcal{C}}(X_i, GT) \\ &\cong \lim_{\mathcal{I}^{\text{opp}}} \operatorname{Hom}_{\mathcal{D}}(FX_i, T), \end{aligned}$$

functorial in T , and hence an isomorphism of functors

$$\operatorname{Hom}_{\mathcal{D}}(F(\operatorname{colim}_{\mathcal{I}} X_i), -) \cong \lim_{\mathcal{I}^{\text{opp}}} \operatorname{Hom}_{\mathcal{D}}(FX_i, -)$$

which by Theorem 8.31 shows that $F(\operatorname{colim}_{\mathcal{I}} X_i)$ is the colimit of the diagram $\mathcal{I} \rightarrow \mathcal{D}$, $i \mapsto FX_i$. \square

The co-theorem states that right adjoints commute with limits:

THEOREM 8.33 (Right adjoints commute with limits). *Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a left adjoint to $G: \mathcal{D} \rightarrow \mathcal{C}$. Let $X: \mathcal{I} \rightarrow \mathcal{D}$ be a functor, and suppose that $\lim_{\mathcal{I}} X$ exists in \mathcal{D} . Then $\lim_{\mathcal{I}}(GX)$ exists and*

$$\lim_{\mathcal{I}}(GX) \cong G(\lim_{\mathcal{I}} X)$$

in \mathcal{C} . \square

EXAMPLE 8.34. Let R be a ring and let A be an (S, R) -bimodule. By Theorem 6.18 the functor

$${}_R\mathbf{Mod} \rightarrow {}_S\mathbf{Mod}, M \mapsto A \otimes_R M$$

is left adjoint (to the functor $\operatorname{Hom}_S(A, -)$), and hence by Theorem 8.32 it commutes with colimits. In particular, it commutes with coproducts:

$$A \otimes_R (\oplus_{i \in I} M_i) = \oplus_{i \in I} (A \otimes_R M_i)$$

and with cokernels:

$$\operatorname{coker}(\operatorname{id} \otimes f: A \otimes_R M \rightarrow A \otimes_R N) = A \otimes_R \operatorname{coker}(f: M \rightarrow N).$$

(Note that the latter gives a one-line proof of Proposition 6.16!).

There is a priori no reason for the functor to commute with limits, and indeed in general $A \otimes_R -$ does not respect kernels (see Exercise 6.13) or (infinite) products (see Exercise 8.11).

EXAMPLE 8.35. The forgetful functor ${}_R\mathbf{Mod} \rightarrow \mathbf{Set}$ is right adjoint to the free module functor (see Example 7.5) and hence by Theorem 8.33 it commutes with limits. For example, this implies that the underlying set of a product of modules is the product of the underlying sets of the modules.

On the other side of the adjunction, we see that the free module functor $I \mapsto R^{(I)}$ must commute with colimits. For example, this contains the (trivial) statement that a basis for the direct sum of free modules is given by the disjoint union of their bases, that is

$$R^{(I)} \oplus R^{(J)} \cong R^{(I \amalg J)}$$

as R -modules.

EXAMPLE 8.36. We have seen in Example 7.6 that the forgetful functor $\mathbf{Top} \rightarrow \mathbf{Set}$ is both right adjoint (to the discrete topology functor) and left adjoint (to the trivial topology functor). It therefore commutes with both limits and colimits, and hence any limit or colimit of topological spaces can be constructed by putting a suitable topology on the limit or colimit of the underlying sets.

Exercises

EXERCISE 8.1. Let X and Y be topological spaces. Show that the cartesian product $X \times Y$ with the product topology is the product of X and Y in the category **Top**.

EXERCISE 8.2. Does the category of pointed topological spaces **Top**_{*} have products and/or coproducts? And if so, what are they? Does the forgetful functor **Top**_{*} \rightarrow **Top** commute with products and/or coproducts?

EXERCISE 8.3. Let \mathcal{C} be a category and let X and Y be objects in \mathcal{C} . Find a category \mathcal{P} in which the products of X and Y are precisely the final objects. Conclude that Proposition 8.2 can be deduced directly from Proposition 4.25.

EXERCISE 8.4. Show that the co-equalizer of the constant maps $0, 1: \{\star\} \rightarrow [0, 1]$ in **Top** is the circle.

EXERCISE 8.5. Let \mathcal{C} be a category. Consider the diagonal functor

$$\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$$

defined by $X \mapsto (X, X)$ and $f \mapsto (f, f)$. When does Δ have a left adjoint? And a right adjoint?

EXERCISE 8.6 (Coproduct of commutative rings). Let R and S be commutative rings.

- (1) Show that there is a unique ring structure on the \mathbf{Z} -module $R \otimes_{\mathbf{Z}} S$ such that

$$(r_1 \otimes s_1)(r_2 \otimes s_2) = r_1 r_2 \otimes s_1 s_2.$$

- (2) Show that $R \rightarrow R \otimes_{\mathbf{Z}} S$, $r \mapsto r \otimes 1$ and $S \rightarrow R \otimes_{\mathbf{Z}} S$, $s \mapsto 1 \otimes s$ are ring homomorphisms.
 (3) Show that $R \otimes_{\mathbf{Z}} S$ is the coproduct of R and S in **CRing**.

EXERCISE 8.7. Let $f: C \rightarrow A$ and $g: C \rightarrow B$ be morphisms in **Ab**. Show that their fibered coproduct exists.

EXERCISE 8.8. Let R be a ring and let

$$\begin{array}{ccc} M & \xrightarrow{f} & P \\ \downarrow g & & \downarrow \varphi \\ Q & \xrightarrow{\psi} & N \end{array}$$

be a commutative square of R -modules.

- (1) Show that M (with the maps f and g) is the pull-back of $P \rightarrow N$ and $Q \rightarrow N$ if and only if the sequence

$$0 \longrightarrow M \xrightarrow{(f,g)} P \oplus Q \xrightarrow{\varphi-\psi} N$$

is exact.

- (2) Show that N (with the maps φ and ψ) is the push-out of $M \rightarrow P$ and $M \rightarrow Q$ if and only if the sequence

$$M \xrightarrow{(f,g)} P \oplus Q \xrightarrow{\varphi-\psi} N \longrightarrow 0$$

is exact.

EXERCISE 8.9. Let $f, g: X \rightarrow Y$ be morphisms in a category \mathcal{C} . Show that the equalizer $\iota: E \rightarrow X$ of f and g is a monomorphism (if it exists).

EXERCISE 8.10. Let \mathcal{C} be a category which has both products and fibered products. Show that \mathcal{C} has equalizers. (Hint: given $f, g: X \rightarrow Y$ consider their fibered product $Z := X \times_Y X$. Show that there are maps $Z \rightarrow X \times X$ and $X \rightarrow X \times X$ whose fibered product is an equalizer of f and g .)

EXERCISE 8.11. Show that the element

$$1 \otimes (1, 1, \dots)$$

of the module $\mathbf{Q} \otimes_{\mathbf{Z}} (\prod_{n>0} \mathbf{Z}/n\mathbf{Z})$ is non-zero. Conclude that the functor $\mathbf{Q} \otimes_{\mathbf{Z}} -$ from \mathbf{Ab} to \mathbf{Ab} does not commute with infinite products.

EXERCISE 8.12. Let $X, Y: \mathcal{I} \rightarrow \mathcal{C}$ be functors, and let $\eta: X \rightarrow Y$ be a morphism of functors. Assume that $\lim_{\mathcal{I}} X$ and $\lim_{\mathcal{I}} Y$ exist. Show that α induces a morphism $\lim_{\mathcal{I}} X \rightarrow \lim_{\mathcal{I}} Y$ in \mathcal{C} . Formulate and prove the analogous statement for colimits.

EXERCISE 8.13. Show that the \lim and colim in Example 8.27 coincide with those described by Propositions 8.28 and 8.29.

EXERCISE 8.14. Let $S_0 \subset S_1 \subset S_2 \cdots$ be an infinite sequence of inclusions of sets. Show that the union $\cup_i S_i$ is the colimit of a suitably chosen diagram $\mathcal{I} \rightarrow \mathbf{Set}$.

EXERCISE 8.15. Let K be a field. Let \mathcal{I} be the category with $\text{ob } \mathcal{I} = \mathbf{N}$ and

$$\text{Hom}_{\mathcal{I}}(i, j) = \begin{cases} \{\star\} & j \leq i \\ \emptyset & j > i \end{cases}$$

Consider the diagram

$$R: \mathcal{I} \rightarrow \mathbf{CRing}, i \mapsto R_i := K[X]/(X^i),$$

(where for $j \leq i$ the map $K[X]/(X^i) \rightarrow K[X]/(X^j)$ is the quotient map). Show that $\lim_{\mathcal{I}} R_i$ exists in \mathbf{CRing} , and is isomorphic to the power series ring $K[[X]]$.

EXERCISE 8.16. Let G be a group and let BG the category of Example 4.7. Let $F: BG \rightarrow \mathbf{Set}$ be a functor.

- (1) Show that $F(\star)$ is a set X equipped with an action of G .
- (2) Show that $\lim F$ is the set of fixed points of the action.
- (3) Show that $\text{colim } F$ is the set of orbits of the action.

EXERCISE 8.17. Let X be a topological space, and let $(U_i)_{i \in I}$ be an open cover. Let \mathcal{I} be the category with $\text{ob } \mathcal{I} = I$ and

$$\text{Hom}(i, j) = \begin{cases} \{\star\} & U_i \subset U_j \\ \emptyset & \text{otherwise} \end{cases}$$

Assume that for all i, j there is a $k \in I$ such that $U_i \cap U_j = U_k$. Show that $\text{colim}_{\mathcal{I}} U_i = X$ in \mathbf{Top} .

EXERCISE 8.18. Let \mathcal{Z} and \mathcal{R} be the categories of Exercise 7.4. Let \mathcal{I} be the category with $\text{ob } \mathcal{I} = \mathbf{N}$ and

$$\text{Hom}_{\mathcal{I}}(i, j) = \begin{cases} \{\star\} & i \leq j \\ \emptyset & i > j \end{cases}$$

Verify that a functor $\mathcal{I} \rightarrow \mathcal{R}$ is the same as an increasing sequence of real numbers

$$x_0 \leq x_1 \leq x_2 \leq \cdots$$

When does this functor have a limit? And a colimit? Verify directly if they are preserved by the left and right adjoints of the inclusion functor $\mathcal{Z} \rightarrow \mathcal{R}$.

EXERCISE 8.19. Prove Proposition 8.29.

EXERCISE 8.20. Show that all limits and colimits exist in the category \mathbf{Top} . (Hint: see Propositions 8.28 and 8.29).

EXERCISE 8.21. Show that \mathbf{Ab} has all limits and colimits. (Hint: for colimits, first do Exercise 8.7).

EXERCISE 8.22. Let R and S be rings, let A be an (R, S) -module and let

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3$$

be an exact sequence of R -modules. Use Theorem 8.33 to prove that the induced sequence

$$0 \longrightarrow \mathrm{Hom}_R(A, M_1) \longrightarrow \mathrm{Hom}_R(A, M_2) \longrightarrow \mathrm{Hom}_R(A, M_3)$$

is exact in ${}_S\mathbf{Mod}$. (See Exercise 2.8 for a more direct approach).

CHAPTER 9

Chain complexes

1. Chain complexes and their homology modules

DEFINITION 9.1. Let R be a ring. A *chain complex* of R -modules consists of

- (1) for every $i \in \mathbf{Z}$ an R -module M_i ,
- (2) for every $i \in \mathbf{Z}$ an R -linear map $d_i: M_i \rightarrow M_{i-1}$

such that for every i the identity $d_i \circ d_{i+1} = 0$ holds.

We depict a chain complex as a diagram

$$\cdots \longrightarrow M_2 \xrightarrow{d_2} M_1 \xrightarrow{d_1} M_0 \xrightarrow{d_0} M_{-1} \longrightarrow \cdots$$

and often denote the chain complex by M_\bullet .

DEFINITION 9.2. A *morphism of chain complexes* $M_\bullet \rightarrow M'_\bullet$ consists of an R -module homomorphism $f_i: M_i \rightarrow M'_i$ for every i , such that the resulting diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & M_2 & \xrightarrow{d_2} & M_1 & \xrightarrow{d_1} & M_0 & \longrightarrow & \cdots \\ & & \downarrow f_2 & & \downarrow f_1 & & \downarrow f_0 & & \\ \cdots & \longrightarrow & M'_2 & \xrightarrow{d'_2} & M'_1 & \xrightarrow{d'_1} & M'_0 & \longrightarrow & \cdots \end{array}$$

commutes. The resulting category of chain complexes of left R -modules is denoted ${}_R\mathbf{Ch}$. The similarly-defined category of chain complexes of right R -modules is denoted \mathbf{Ch}_R .

The condition $d_i \circ d_{i+1} = 0$ in a chain complex

$$\cdots \longrightarrow M_{i+1} \xrightarrow{d_{i+1}} M_i \xrightarrow{d_i} M_{i-1} \longrightarrow \cdots$$

implies that $\text{im } d_{i+1} \subset \ker d_i$ inside M_i .

DEFINITION 9.3. Let M_\bullet be a chain complex of R -modules and i an integer. The i -th homology module of M_\bullet is the R -module

$$H_i(M_\bullet) := \frac{\ker d_i}{\operatorname{im} d_{i+1}}.$$

If $f: M_\bullet \rightarrow M'_\bullet$ is a morphism of chain complexes, then $f_i: M_i \rightarrow M'_i$ induces a morphism

$$H_i(f): H_i(M_\bullet) \rightarrow H_i(M'_\bullet)$$

(see Exercise 9.1). We obtain for every i a functor

$$H_i: {}_R\mathbf{Ch} \rightarrow {}_R\mathbf{Mod}.$$

These modules measure the failure of the sequence M_\bullet to be exact, in the following sense.

LEMMA 9.4. *A chain complex*

$$\cdots \longrightarrow M_2 \xrightarrow{d_2} M_1 \xrightarrow{d_1} M_0 \xrightarrow{d_0} M_{-1} \longrightarrow \cdots$$

is exact if and only if $H_i(M_\bullet) = 0$ for all $i \in \mathbf{Z}$. □

2. The long exact sequence

Let M_\bullet , N_\bullet and P_\bullet be chain complexes of R -modules. We say that a sequence of morphisms

$$0 \longrightarrow M_\bullet \xrightarrow{\alpha} N_\bullet \xrightarrow{\beta} P_\bullet \longrightarrow 0$$

is a short exact sequence in ${}_R\mathbf{Ch}$ if it is termwise exact. In other words, if in the commutative diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & M_{i+1} & \longrightarrow & M_i & \longrightarrow & M_{i-1} \longrightarrow \cdots \\ & & \downarrow \alpha_{i+1} & & \downarrow \alpha_i & & \downarrow \beta_{i-1} \\ \cdots & \longrightarrow & N_{i+1} & \longrightarrow & N_i & \longrightarrow & N_{i-1} \longrightarrow \cdots \\ & & \downarrow \beta_{i+1} & & \downarrow \beta_i & & \downarrow \beta_{i-1} \\ \cdots & \longrightarrow & P_{i+1} & \longrightarrow & P_i & \longrightarrow & P_{i-1} \longrightarrow \cdots \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

all the columns are short exact sequences of R -modules.

THEOREM 9.5. *Let*

$$0 \longrightarrow M_{\bullet} \xrightarrow{\alpha} N_{\bullet} \xrightarrow{\beta} P_{\bullet} \longrightarrow 0$$

be a short exact sequence of chain complexes of R -modules. Then there exists an exact sequence

$$\begin{array}{ccccccc} & & \cdots & \longrightarrow & H_{i+1}(N_{\bullet}) & \xrightarrow{H_{i+1}(\beta)} & H_{i+1}(P_{\bullet}) & \longrightarrow & \\ & & & & \searrow & & \swarrow & & \\ \hookrightarrow & H_i(M_{\bullet}) & \xrightarrow{H_i(\alpha)} & H_i(N_{\bullet}) & \xrightarrow{H_i(\beta)} & H_i(P_{\bullet}) & \longrightarrow & & \\ & & & \swarrow & & \searrow & & & \\ \hookrightarrow & H_{i-1}(M_{\bullet}) & \xrightarrow{H_{i-1}(\alpha)} & H_{i-1}(N_{\bullet}) & \longrightarrow & \cdots & & & \end{array}$$

of R -modules.

The resulting exact sequence of homology modules is called the *long exact sequence of homology* associated to the short exact sequence of chain complexes.

PROOF OF THEOREM 9.5. We omit the proof, which is a tedious diagram chase in the style of the proof of the Snake Lemma (Theorem 2.2). In fact, the Snake Lemma is a special case of the theorem, obtained by assuming the M_i , N_i and P_i vanish for $i \notin \{0, 1\}$. \square

3. The homotopy category

DEFINITION 9.6. Let R be a ring and let $f: M_{\bullet} \rightarrow M'_{\bullet}$ and $g: M_{\bullet} \rightarrow M'_{\bullet}$ be morphisms of chain complexes of R -modules. A *homotopy* from f to g consists of a collection of R -linear maps $h_i: M_i \rightarrow M'_{i+1}$ indexed by $i \in \mathbf{Z}$, such that for every $i \in \mathbf{Z}$ the identity

$$g_i - f_i = d'_{i+1}h_i + h_{i-1}d_i$$

holds in $\text{Hom}_R(M_i, M'_i)$. We say that f and g are *homotopic*, and write $f \sim g$, if there exists a homotopy from f to g .

It is convenient to keep track of these maps in a diagram:

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & M_2 & \xrightarrow{d_2} & M_1 & \xrightarrow{d_1} & M_0 & \longrightarrow & \cdots \\
 & & \downarrow f_2 & \parallel g_2 & \downarrow f_1 & \parallel g_1 & \downarrow f_0 & \parallel g_0 & \\
 & \swarrow & & & \swarrow & & \swarrow & & \\
 \cdots & \longrightarrow & M'_2 & \xrightarrow{d'_2} & M'_1 & \xrightarrow{d'_1} & M'_0 & \longrightarrow & \cdots
 \end{array}$$

but note that in the definition of homotopy it is *not* required that the h_i 's commute with the horizontal maps d in any way.

The equation defining homotopy can be remembered as

$$g - f = dh + hd,$$

omitting the indices which can be reinserted in only one meaningful way.

PROPOSITION 9.7. *Let R be a ring and M_\bullet, M'_\bullet chain complex of R -modules. Then homotopy is an equivalence relation on the set $\text{Hom}_R \mathbf{Ch}(M_\bullet, M'_\bullet)$. \square*

PROPOSITION 9.8. *Let R be a ring and $f, g: M_\bullet \rightarrow M'_\bullet$ homotopic morphisms of chain complexes of R -modules. Then*

- (1) *for any morphism $s: M'_\bullet \rightarrow N_\bullet$ in ${}_R \mathbf{Ch}$, the compositions sf and sg are homotopic;*
- (2) *for any morphism $t: N_\bullet \rightarrow M_\bullet$ in ${}_R \mathbf{Ch}$, the compositions ft and gt are homotopic.*

PROOF. Let $(h_i)_i$ be a homotopy from f to g . In the first case, one verifies that $(s_{i+1}h_i)_i$ is a homotopy from sf to sg , and in the second case, that $(h_it_i)_i$ is a homotopy from ft to gt . \square

DEFINITION 9.9. The *homotopy category of chain complexes of R -modules*, denoted ${}_R \mathbf{Ho}$, is the category with

- (1) $\text{ob } {}_R \mathbf{Ho} := \text{ob } {}_R \mathbf{Ch}$
- (2) $\text{Hom}_{{}_R \mathbf{Ho}}(M_\bullet, N_\bullet) := \text{Hom}_{{}_R \mathbf{Ch}}(M_\bullet, N_\bullet) / \sim$

where composition and identity maps are inherited from composition and identity maps in ${}_R \mathbf{Ch}$.

Proposition 9.8 guarantees that composition in ${}_R \mathbf{Ho}$ is well-defined.

PROPOSITION 9.10. *Let $f, g: M_\bullet \rightarrow M'_\bullet$ be homotopic maps in \mathbf{Ch}_R , and let i be an integer. Then $H_i(f) = H_i(g)$ as maps $H_i(M_\bullet) \rightarrow H_i(M'_\bullet)$.*

PROOF. By definition, an element of $H_i(M_\bullet)$ is a coset

$$\bar{x} := x + \text{im}(d_{i+1})$$

for some $x \in \ker(d_i)$. We have

$$\begin{aligned} H_i(f)(\bar{x}) - H_i(g)(\bar{x}) &= f(x) - g(x) + \text{im}(d'_{i+1}) \\ &= d'_{i+1}(h_i(x)) + h_{i-1}(d_i(x)) + \text{im}(d'_{i+1}). \end{aligned}$$

Since $d'_{i+1}(h_i(x)) \in \text{im}(d'_{i+1})$, the first term vanishes in $H_i(M'_\bullet)$. Since $x \in \ker(d_i)$, also the second term vanishes, and $H_i(f) = H_i(g)$. \square

A consequence of Proposition 9.10 is that the functors H_i on $R\mathbf{Ch}$ induce functors

$$H_i: R\mathbf{Ho} \rightarrow R\mathbf{Mod}, M_\bullet \mapsto H_i(M_\bullet)$$

on the homotopy category of chain complexes.

REMARK 9.11. As the terminology suggests, there is a close relationship with the notions of homotopic maps and homology groups in algebraic topology. Homology groups of topological spaces are usually defined in terms of the functor

$$C: \mathbf{Top} \rightarrow \mathbf{zCh}, X \mapsto C_\bullet(X)$$

that maps a space X to the chain complex of *singular chains*. One then defines the i -th homology group of X by

$$H_i(X, \mathbf{Z}) := H_i(C_\bullet(X))$$

and obtains functors $H_i: \mathbf{Top} \rightarrow \mathbf{Ab}$.

If $f, g: X \rightarrow Y$ are homotopic continuous maps, then their induced maps $C_\bullet(X) \rightarrow C_\bullet(Y)$ are homotopic maps of chain complexes, and hence their induced maps $H_i(X, \mathbf{Z}) \rightarrow H_i(Y, \mathbf{Z})$ coincide.

REMARK 9.12. Chain complexes form a mathematical context in which three layers play a role: objects (chain complexes), morphisms (morphisms of chain complexes), and maps between morphisms (homotopies). See Remark 5.20 for two other such contexts: topological spaces (spaces, continuous maps, homotopies), and categories (categories, functors, morphisms of functors).

Exercises

EXERCISE 9.1 (Functoriality of homology). Let $f: M_\bullet \rightarrow M'_\bullet$ be a morphism of chain complexes of R -modules.

- (1) Show $f_i(\ker d_i) \subset \ker d'_i$;
- (2) Show $f_i(\operatorname{im} d_{i+1}) \subset \operatorname{im} d'_{i+1}$;
- (3) Conclude that f_i induces an R -linear map $H_i(M) \rightarrow H_i(M')$.

EXERCISE 9.2. Let R be a non-zero ring. Show that the two chain complexes

$$\begin{array}{ccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \cdots \\ \cdots & \longrightarrow & 0 & \longrightarrow & R & \xrightarrow{\operatorname{id}} & R & \longrightarrow & 0 & \longrightarrow & \cdots \end{array}$$

are isomorphic in ${}_R\mathbf{Ho}$.

EXERCISE 9.3. Let $n > 1$ and let f be the morphism in \mathbf{zCh} given by the diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & \mathbf{Z} & \xrightarrow{n} & \mathbf{Z} & \longrightarrow & 0 & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \downarrow \pi & & \downarrow & & \\ \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \mathbf{Z}/n\mathbf{Z} & \longrightarrow & 0 & \longrightarrow & \cdots \end{array}$$

with π the canonical map. Show that $H_i(f)$ is an isomorphism for all i , but that f is not an isomorphism in \mathbf{zHo} .

EXERCISE 9.4. Let R and S be rings. A functor $F: {}_R\mathbf{Mod} \rightarrow {}_S\mathbf{Mod}$ is called *additive* if for all R -modules M, N the map

$$F: \operatorname{Hom}_R(M, N) \rightarrow \operatorname{Hom}_S(FM, FN)$$

is a homomorphism of abelian groups. Let A be an (R, S) -bimodule. Show that the functor

$${}_R\mathbf{Mod} \rightarrow {}_S\mathbf{Mod}, M \mapsto \operatorname{Hom}_R(A, M)$$

is additive.

EXERCISE 9.5. Show that an additive functor $F: {}_R\mathbf{Mod} \rightarrow {}_S\mathbf{Mod}$ induces functors ${}_R\mathbf{Ch} \rightarrow {}_S\mathbf{Ch}$ and ${}_R\mathbf{Ho} \rightarrow {}_S\mathbf{Ho}$.

EXERCISE 9.6. Let R and S be rings and A an (R, S) -bimodule. For a chain complex M_\bullet in ${}_R\mathbf{Ch}$ define a chain complex M'_\bullet in \mathbf{Ch}_S by

- (1) $M'_i := \operatorname{Hom}_R(M_{-i}, A)$

(2) $d_i: M'_i \rightarrow M'_{i-1}$ the map induced from $d_{1-i}: M_{1-i} \rightarrow M_{-i}$

Verify that M'_\bullet is a chain complex of right S -modules, and that the operation $M_\bullet \mapsto M'_\bullet$ defines functors ${}_R\mathbf{Ch}^{\text{opp}} \rightarrow \mathbf{Ch}_S$ and ${}_R\mathbf{Ho}^{\text{opp}} \rightarrow \mathbf{Ho}_S$.

EXERCISE 9.7 (\star). Let K be a field. Let $\prod_{i \in \mathbf{Z}} \mathbf{Vec}_K$ be the category whose objects are sequences $(V_i)_{i \in \mathbf{Z}}$ of K -vector spaces, and whose morphisms are sequences $(f_i)_{i \in \mathbf{Z}}$ of K -linear maps. Show that the functor

$$\mathbf{Ho}_K \rightarrow \prod_{i \in \mathbf{Z}} \mathbf{Vec}_K, V_\bullet \mapsto (H_i(V_\bullet))_i$$

is an equivalence of categories.

CHAPTER 10

Free resolutions

1. Definition and existence

DEFINITION 10.1. Let R be a ring and let M be an R -module. A *free resolution* of M is an exact sequence of R -modules

$$\cdots \longrightarrow F_2 \xrightarrow{d_2} F_1 \xrightarrow{d_1} F_0 \xrightarrow{\pi} M \longrightarrow 0$$

in which the modules F_i are free.

EXAMPLE 10.2. If M itself is free, then the exact sequence

$$\cdots \longrightarrow 0 \longrightarrow 0 \longrightarrow M \xrightarrow{\text{id}} M \longrightarrow 0$$

is a free resolution (with $F_0 = M$ and $F_i = 0$ for $i \neq 0$). We will usually suppress leading zeroes from the notation, and simply write

$$0 \longrightarrow M \longrightarrow M \longrightarrow 0$$

for the above resolution.

EXAMPLE 10.3. Let R be a domain. If $I \subset R$ is a non-zero principal ideal, then I is free of rank 1 as an R -module (see Exercise 1.6), hence the exact sequence

$$0 \longrightarrow I \longrightarrow R \longrightarrow R/I \longrightarrow 0$$

is a free resolution of the R -module R/I .

EXAMPLE 10.4. If R is a principal ideal domain, and M a finitely generated R -module, then we have seen in Corollary 3.2 that M has a free resolution of the form

$$0 \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$$

with F_0 and F_1 free R -modules of finite rank.

EXAMPLE 10.5. Let K be a field, and let $R = K[X, Y]$. Then the sequence

$$0 \longrightarrow R \xrightarrow{d_2} R \oplus R \xrightarrow{d_1} R \xrightarrow{\pi} K[X, Y]/(X, Y) \longrightarrow 0$$

with $d_2(1) = (Y, -X)$, $d_1(1, 0) = X$ and $d_1(0, 1) = Y$, and π the quotient map is a free resolution of the R -module $K[X, Y]/(X, Y)$.

Every R -module has a free resolution, although in general the resolution will be infinite, and moreover the free modules occurring in it will not be of finite rank.

PROPOSITION 10.6. *Let R be a ring and let M be an R -module. Then there exists a free resolution of M .*

PROOF. Choose a generating set I of M , and let $F_0 = R^{(I)}$ be the free R -module with basis I . Then we have a natural surjective map $\pi: F_0 \rightarrow M$, and hence an exact sequence

$$F_0 \xrightarrow{\pi} M \longrightarrow 0.$$

Now choose a generating set I_1 of the R -module $K_0 := \ker \pi \subset F_0$ and let $F_1 = R^{(I_1)}$. The natural map $F_1 \rightarrow K_0 \subset F_0$ extends the above to an exact sequence

$$F_1 \xrightarrow{d_1} F_0 \xrightarrow{\pi} M \longrightarrow 0.$$

Repeating this argument with $K_1 := \ker d_1$ and so forth, we obtain an exact sequence

$$\cdots \longrightarrow F_2 \xrightarrow{d_2} F_1 \xrightarrow{d_1} F_0 \xrightarrow{\pi} M \longrightarrow 0,$$

with the F_i free, as we had to show. \square

REMARK 10.7. From a free resolution

$$\cdots \longrightarrow F_2 \xrightarrow{d_2} F_1 \xrightarrow{d_1} F_0 \xrightarrow{\pi} M \longrightarrow 0$$

of M we obtain a chain complex F_\bullet of the form

$$\cdots \longrightarrow F_2 \xrightarrow{d_2} F_1 \xrightarrow{d_1} F_0 \longrightarrow 0 \longrightarrow 0 \longrightarrow \cdots$$

by setting set $F_i = 0$ for $i < 0$. This chain complex satisfies

$$H_i(F_\bullet) \cong \begin{cases} M & i = 0 \\ 0 & i \neq 0 \end{cases}$$

Conversely, given a pair (F_\bullet, α) consisting of

- (1) a chain complex F_\bullet

(2) an isomorphism $\alpha: H_0(F_\bullet) \rightarrow M$

with F_i free for all i and zero for $i < 0$, and with $H_i(F_\bullet) = 0$ for all $i \neq 0$, the sequence

$$\cdots \longrightarrow F_2 \longrightarrow F_1 \longrightarrow F_0 \xrightarrow{\pi} M \longrightarrow 0,$$

where π is induced by α , is a free resolution of M .

It will often be convenient to think of a free resolution as a pair (F_\bullet, α) consisting of a chain complex F_\bullet and an isomorphism α as above.

2. The free resolution functor

THEOREM 10.8. *Let M and M' be R -modules. Let F_\bullet and F'_\bullet be free resolutions of M and M' respectively. Let $\varphi: M \rightarrow M'$ be a morphism of R -modules. Then there exists a morphism $f: F_\bullet \rightarrow F'_\bullet$ such that $H_0(f) = \varphi$. Moreover, f is unique up to homotopy.*

PROOF. The existence of f amounts to the existence of R -linear maps f_i making the diagram (with exact rows)

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & F_2 & \xrightarrow{d_2} & F_1 & \xrightarrow{d_1} & F_0 & \xrightarrow{\pi} & M & \longrightarrow & 0 \\ & & \downarrow f_2 & & \downarrow f_1 & & \downarrow f_0 & & \downarrow \varphi & & \\ \cdots & \longrightarrow & F'_2 & \xrightarrow{d'_2} & F'_1 & \xrightarrow{d'_1} & F'_0 & \xrightarrow{\pi'} & M' & \longrightarrow & 0 \end{array}$$

commute. We will construct such a diagram inductively.

Let $S_0 \subset F_0$ be a basis of the free module F_0 . For every $s \in S_0$, choose an $s' \in F'_0$ such that $\pi'(s') = \varphi\pi(s)$. Such s' exists by the surjectivity of π' . Now, since F_0 is free with basis S_0 , there exists a unique R -linear map $f_0: F_0 \rightarrow F'_0$ that maps every $s \in S_0$ to its chosen counterpart $s' \in F'_0$. By construction the right-hand square commutes.

Next, let $S_1 \subset F_1$ be a basis of F_1 . For every $s \in S_1$, we have $\pi d_1(s) = 0$ by the exactness of the top-row, hence $\pi' f_0 d_1(s) = 0$ by the commutativity of the right-hand square, and hence $f_0 d_1(s) \in \ker \pi'$. We conclude that there exists an element $s' \in F'_1$ with $d'_1(s') = f_0 d_1(s)$. Choosing such an s' for every $s \in S_1$ yields an R -linear map $f_1: F_1 \rightarrow F'_1$ as above. Repeating the argument, we construct maps f_i as required.

For the uniqueness assertion in the theorem, assume that we have chain complex homomorphisms $f: F_\bullet \rightarrow F'_\bullet$ and $g: F_\bullet \rightarrow F'_\bullet$ with $H_0(f) = H_0(g) = \varphi$. Let $\delta := g - f$. We need to show that there

are h_i as in the diagram below

$$\begin{array}{ccccccccccc}
 \cdots & \longrightarrow & F_2 & \xrightarrow{d_2} & F_1 & \xrightarrow{d_1} & F_0 & \xrightarrow{\pi} & M & \longrightarrow & 0 \\
 & & \downarrow \delta_2 & \swarrow h_1 & \downarrow \delta_1 & \swarrow h_0 & \downarrow \delta_0 & & \downarrow 0 & & \\
 \cdots & \longrightarrow & F'_2 & \xrightarrow{d'_2} & F'_1 & \xrightarrow{d'_1} & F'_0 & \xrightarrow{\pi'} & M' & \longrightarrow & 0
 \end{array}$$

satisfying

$$\begin{aligned}
 \delta_0 &= d'_1 h_0 \\
 \delta_i &= d'_{i+1} h_i + h_{i-1} d_i \quad (i \geq 1)
 \end{aligned}$$

As in the proof of the first part of the theorem, we construct these h_i inductively.

For the base step, choose a basis S_0 of F_0 and for every $s \in S_0$ choose an $s' \in F'_1$ with $d'_1 s' = \delta_0 s$. Such s' exists, since $\pi' \delta_0 s = 0$, and the bottom row in the diagram is exact. Since F_0 is free with basis S_0 , there is a (unique) morphism $h_0: F_0 \rightarrow F'_1$ with $s \rightarrow s'$ for every $s \in S_0$, and we have $\delta_0 = d'_1 h_0$ by construction.

Next, let S_1 be a basis of F_1 , and choose for every $s \in S_1$ an $s' \in F'_2$ with $d'_2 s' = \delta_1 s - h_0 d_1 s$. Such s' exists, since

$$d'_1(\delta_1 s - h_0 d_1 s) = d'_1 \delta_1 s - d'_1 h_0 d_1 s = \delta_0 d_1 s - \delta_0 d_1 s = 0.$$

As before, the collection of s' defines a map $h_1: F_1 \rightarrow F'_2$, and repeating the argument gives a collection of maps h_i defining the desired homotopy between f and g . \square

COROLLARY 10.9. *If F_\bullet and F'_\bullet are free resolutions of M , then F_\bullet and F'_\bullet are isomorphic in ${}_R\mathbf{Ho}$.*

PROOF. The proof is ‘abstract nonsense’ and quite similar to the argument that showed that final objects are unique up to unique isomorphism (see Proposition 4.25).

Take $M' := M$ and apply Theorem 10.8 to $\text{id}: M \rightarrow M'$ and $\text{id}: M' \rightarrow M$ to obtain morphisms $f: F_\bullet \rightarrow F'_\bullet$ and $g: F'_\bullet \rightarrow F_\bullet$. Then apply Theorem 10.8 again to id_M and $\text{id}_{M'}$ to show that gf is homotopic to id_{F_\bullet} and fg is homotopic to $\text{id}_{F'_\bullet}$. This gives equalities $fg = \text{id}_{F'_\bullet}$ and $gf = \text{id}_{F_\bullet}$ in ${}_R\mathbf{Ho}$, which shows that f and g are mutually inverse isomorphisms in ${}_R\mathbf{Ho}$. \square

EXAMPLE 10.10. The zero module $M = \{0\}$ has the zero resolution, but also the non-trivial free resolution

$$0 \longrightarrow R \xrightarrow{\text{id}} R \xrightarrow{\pi} M \longrightarrow 0,$$

hence the corresponding complexes

$$\begin{aligned} \cdots &\longrightarrow 0 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0 \longrightarrow \cdots \\ \cdots &\longrightarrow 0 \longrightarrow R \longrightarrow R \longrightarrow 0 \longrightarrow \cdots \end{aligned}$$

are isomorphic in ${}_R\mathbf{Ho}$. See also Exercise 9.2.

We can now summarise this section into one powerful theorem.

THEOREM 10.11. *There exists a functor*

$$F: {}_R\mathbf{Mod} \rightarrow {}_R\mathbf{Ho}, M \mapsto F_\bullet(M)$$

and an isomorphism of functors

$$\alpha: \mathbf{H}_0 \circ F \xrightarrow{\sim} \text{id}_{{}_R\mathbf{Mod}}$$

such that for every R -module M , the complex $F_\bullet(M)$ together with the isomorphism α_M forms a free resolution of M .

The proof goes directly against our basic principle that ‘constructions depending on choices do not give rise to functors’.

PROOF OF THEOREM 10.11. Using Proposition 10.6, choose for every R -module M a free resolution

$$\cdots \longrightarrow F_2(M) \longrightarrow F_1(M) \longrightarrow F_0(M) \xrightarrow{\pi_M} M \longrightarrow 0.$$

This defines for every R -module M an object $F_\bullet(M) \in {}_R\mathbf{Ho}$, and an isomorphism $\alpha_M: \mathbf{H}_0(F_\bullet(M)) \xrightarrow{\sim} M$ (induced by π_M).

Now for every $\varphi: M \rightarrow N$ in ${}_R\mathbf{Mod}$, Theorem 10.8 gives a *unique* morphism $F_\bullet(\varphi): F_\bullet(M) \rightarrow F_\bullet(N)$ such that the square of R -modules

$$\begin{array}{ccc} \mathbf{H}_0(F_\bullet(M)) & \xrightarrow{\alpha_M} & M \\ \downarrow H_0(F_\bullet(\varphi)) & & \downarrow \varphi \\ \mathbf{H}_0(F_\bullet(N)) & \xrightarrow{\alpha_N} & N \end{array}$$

commutes. This provides the necessary data for a functor F_\bullet , and immediately shows that α is an isomorphism of functors, provided that the data defining F_\bullet indeed forms a functor.

For this, we need to check that F_\bullet respects identity and composition. But this follows quite formally from the uniqueness statement in Theorem 10.8. Given an R -module M , both $\text{id}_{F_\bullet(M)}$ and $F_\bullet(\text{id})$ induce the identity on $H_0(F_\bullet(M)) = M$, so they must be homotopic and hence they define the same morphism in ${}_R\mathbf{Ho}$. Similarly, given $f: M \rightarrow N$ and $g: N \rightarrow P$ then both

$$F_\bullet(g)F_\bullet(f): F_\bullet(M) \rightarrow F_\bullet(N)$$

and

$$F_\bullet(gf): F_\bullet(M) \rightarrow F_\bullet(N)$$

induce the map $gf: M \rightarrow P$ on H_0 , so they must be homotopic and hence define the same morphism in ${}_R\mathbf{Ho}$. \square

Exercises

EXERCISE 10.1. Let K be a field and $R = K[X, Y]$. Show that the R -module $M = K[X, Y]/(X, Y)$ does not have a free resolution of the form

$$0 \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0.$$

EXERCISE 10.2. Let K be a field and consider the subring $R = K[X^2, X^3]$ of the polynomial ring $K[X]$. Let M be the R -module $R/(X^2, X^3)$. Find a free resolution of M .

EXERCISE 10.3. Let R be a commutative ring.

- (1) Assume that $r \in R$ is not a zero divisor in R . Show that R/rR has a free resolution of the form

$$0 \longrightarrow R \longrightarrow R \longrightarrow R/(r) \longrightarrow 0.$$

- (2) Let $r, s \in R$. Assume that r is not a zero divisor in R , and that \bar{s} is not a zero divisor in R/rR . Show that $R/(s, r)$ has a free resolution of the form

$$0 \longrightarrow R \longrightarrow R^2 \longrightarrow R \longrightarrow R/(r, s) \longrightarrow 0.$$

- (3, \star) Try to formulate and prove an analogous statement for modules of the form $R/(r, s, t)$, etcetera.

EXERCISE 10.4. Let n be a positive integer, and consider the ring $R := \mathbf{Z}[X]/(X^n - 1)$. Let M be the quotient module $R/(X - 1)$, $\pi: R \rightarrow M$ the quotient map. Show that

$$\cdots \xrightarrow{\beta} R \xrightarrow{\alpha} R \xrightarrow{\beta} R \xrightarrow{\alpha} R \xrightarrow{\pi} M \longrightarrow 0,$$

with $\alpha(r) = (X - 1)r$ and $\beta(r) = (X^{n-1} + \cdots + X + 1)r$, is a free resolution of the R -module M .

EXERCISE 10.5. Let $R = \mathbf{Z}/4\mathbf{Z}$ and let M be the R -module $\mathbf{Z}/2\mathbf{Z}$. Find a free resolution of M .

EXERCISE 10.6. Let K be a field, $n > 1$ and let R be the matrix ring $\text{Mat}_n(K)$. Let $M = K^n$ be the left R -module of column vectors. Show that M does not have a finite free resolution consisting of finitely generated free R -modules.

EXERCISE 10.7. Let $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ be a short exact sequence of R -modules. Show that there exist free R -modules F_1, F_2, F_3 , and a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & F_1 & \longrightarrow & F_2 & \longrightarrow & F_3 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & M_1 & \longrightarrow & M_2 & \longrightarrow & M_3 & \longrightarrow & 0 \end{array}$$

with exact rows and surjective vertical maps.

EXERCISE 10.8 (Free resolution of a short exact sequence). Let $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ be a short exact sequence of R -modules. Show that there exist free resolutions

$$\cdots \rightarrow F_{i,2} \rightarrow F_{i,1} \rightarrow F_{i,0} \rightarrow M_i \rightarrow 0$$

for $i = 1, 2, 3$, and a short exact sequence

$$0 \rightarrow F_{1,\bullet} \rightarrow F_{2,\bullet} \rightarrow F_{3,\bullet} \rightarrow 0$$

of chain complexes compatible with the exact sequence $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$.

EXERCISE 10.9 (Uniqueness of free resolution functor). Let F and G be functors ${}_R\mathbf{Mod} \rightarrow {}_R\mathbf{Ho}$. Let $\alpha: H_0 \circ F \xrightarrow{\sim} \text{id}$ and $\beta: H_0 \circ G \xrightarrow{\sim} \text{id}$ be isomorphisms. Assume that for every M the pairs $(F(M), \alpha)$ and $(G(M), \beta)$ are free resolutions of M . Show that the functors F and G are isomorphic.

EXERCISE 10.10. Let M_\bullet be a chain complex of R -modules with $M_i = 0$ for all $i \neq 0, 1$. Show that there exists a chain complex F_\bullet and a morphism $\alpha: F_\bullet \rightarrow M_\bullet$ such that

- (1) $H_i(\alpha)$ is an isomorphism for all i , and
- (2) F_i is free for all i .

EXERCISE 10.11 (\star). Let M_\bullet be a chain complex of R -modules with $M_i = 0$ for all $i < 0$. Show that there exists an F_\bullet and α as in Exercise 10.10.

CHAPTER 11

The Ext functors

1. The functors Ext^n

Let R be a ring. If M_\bullet is the complex of R -modules given by

$$\cdots \longrightarrow M_{i+1} \longrightarrow M_i \longrightarrow M_{i-1} \longrightarrow \cdots$$

and if N is an R -module, then the induced sequence

$$\cdots \longrightarrow \text{Hom}_R(M_{i-1}, N) \longrightarrow \text{Hom}_R(M_i, N) \longrightarrow \text{Hom}_R(M_{i+1}, N) \longrightarrow \cdots$$

forms a complex of abelian groups, with the group $\text{Hom}_R(M_{-i}, N)$ in degree i . This determines a functor

$${}_R\mathbf{Ch}^{\text{opp}} \times {}_R\mathbf{Mod} \rightarrow \mathbf{zCh}$$

and it induces a functor

$${}_R\mathbf{Ho}^{\text{opp}} \times {}_R\mathbf{Mod} \rightarrow \mathbf{zHo}.$$

on the homotopy categories, by Exercise 9.6.

DEFINITION 11.1. Let n be an integer. We define the functor

$$\text{Ext}_R^n(-, -): {}_R\mathbf{Mod}^{\text{opp}} \times {}_R\mathbf{Mod} \rightarrow \mathbf{Ab}, (M, N) \mapsto \text{Ext}_R^n(M, N)$$

as the composition of the following functors:

- (1) the free resolution functor of Theorem 10.11 (applied to the first coordinate)

$$F_\bullet(-) \times \text{id}: {}_R\mathbf{Mod}^{\text{opp}} \times {}_R\mathbf{Mod} \longrightarrow {}_R\mathbf{Ho}^{\text{opp}} \times {}_R\mathbf{Mod}$$

- (2) the functor induced by Hom_R as above

$${}_R\mathbf{Ho}^{\text{opp}} \times {}_R\mathbf{Mod} \longrightarrow \mathbf{zHo}$$

- (3) the homology functor

$$H_{-n}: \mathbf{zHo} \rightarrow \mathbf{Ab},$$

which is well-defined by Proposition 9.10.

In other words, if

$$\cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$$

is a free resolution of M , then the group $\text{Ext}_R^n(M, N)$ is defined as the quotient group

$$\text{Ext}_R^n(M, N) = \frac{\ker(\text{Hom}_R(F_n, N) \rightarrow \text{Hom}_R(F_{n+1}, N))}{\text{im}(\text{Hom}_R(F_{n-1}, N) \rightarrow \text{Hom}_R(F_n, N))},$$

where the maps between the Hom groups are induced from the maps in the free resolution, and where we set $F_i = 0$ for $i < 0$, as before.

REMARK 11.2. A priori the functor $\text{Ext}_R^n(-, -)$ depends on the choices of free resolutions involved in $F_\bullet(-)$, but different choices give rise to isomorphic functors $\text{Ext}_R^n(-, -)$.

EXAMPLE 11.3. Let m be a positive integer. We compute the groups $\text{Ext}_{\mathbf{Z}}^n(\mathbf{Z}/m\mathbf{Z}, \mathbf{Z})$ using the definition. As a first step we need to find a free resolution of $\mathbf{Z}/m\mathbf{Z}$. The obvious choice leads to the complex

$$F_\bullet = [\cdots \longrightarrow 0 \longrightarrow \mathbf{Z} \xrightarrow{m} \mathbf{Z} \longrightarrow 0 \longrightarrow 0 \longrightarrow \cdots]$$

with $F_1 = F_0 = \mathbf{Z}$. Note that $\text{Hom}(\mathbf{Z}, \mathbf{Z}) = \mathbf{Z}$, so that applying the contravariant additive functor $\text{Hom}_{\mathbf{Z}}(-, \mathbf{Z})$ to this complex gives a complex of the form

$$H_\bullet = [\cdots \longrightarrow 0 \longrightarrow 0 \longrightarrow \mathbf{Z} \longrightarrow \mathbf{Z} \longrightarrow 0 \longrightarrow \cdots]$$

with $H_0 = H_{-1} = \mathbf{Z}$. One checks that the map $\mathbf{Z} \rightarrow \mathbf{Z}$ is multiplication by m . We find $H_i(H_\bullet) = 0$ for all $i \neq -1$, and $H_{-1}(H_\bullet) = \mathbf{Z}/m\mathbf{Z}$. From this we conclude

$$\text{Ext}_{\mathbf{Z}}^n(\mathbf{Z}/m\mathbf{Z}, \mathbf{Z}) \cong \begin{cases} \mathbf{Z}/m\mathbf{Z} & n = 1 \\ 0 & n \neq 1 \end{cases}$$

By construction, the groups $\text{Ext}^n(M, N)$ are zero for $n < 0$.

PROPOSITION 11.4. *The functors $\text{Ext}_R^0(-, -)$ and $\text{Hom}_R(-, -)$ are isomorphic.*

PROOF. Let

$$\cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$$

be a free resolution of an R -module M . Then by Exercise 2.9 we get an exact sequence

$$0 \longrightarrow \text{Hom}(M, N) \longrightarrow \text{Hom}(F_0, N) \longrightarrow \text{Hom}(F_1, N).$$

Using the definition of Ext^0 we find

$\text{Ext}^0(M, N) = H_0(\text{Hom}(F_\bullet, N)) = \ker(\text{Hom}(F_0, N) \rightarrow \text{Hom}(F_1, N))$,
 which gives an isomorphism $\text{Ext}^0(M, N) = \text{Hom}(M, N)$, functorial in M . \square

We will see that the module $\text{Ext}^1(M, N)$ is in bijection with isomorphism classes of short exact sequences

$$0 \rightarrow N \rightarrow E \rightarrow M \rightarrow 0$$

of R -modules. There is also an interpretation to the modules $\text{Ext}^n(M, N)$ with $n > 1$ in terms of exact sequences

$$0 \rightarrow N \rightarrow E_1 \rightarrow \cdots \rightarrow E_n \rightarrow M \rightarrow 0,$$

but the statement is more delicate.

2. The long exact sequence

THEOREM 11.5. *Let N be an R -module, and let*

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$$

be a short exact sequence of R -modules. Then there is a natural exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}(M_3, N) & \longrightarrow & \text{Hom}(M_2, N) & \longrightarrow & \text{Hom}(M_1, N) \\ & & \searrow & & \searrow & & \searrow \\ & & \text{Ext}^1(M_3, N) & \longrightarrow & \text{Ext}^1(M_2, N) & \longrightarrow & \text{Ext}^1(M_1, N) \\ & & \searrow & & \searrow & & \searrow \\ & & \text{Ext}^2(M_3, N) & \longrightarrow & \text{Ext}^2(M_2, N) & \longrightarrow & \cdots \end{array}$$

of abelian groups.

PROOF. Choose free resolutions of the M_i as in Exercise 10.8, so that we have a short exact sequence of complexes

$$0 \rightarrow F_{1,\bullet} \rightarrow F_{2,\bullet} \rightarrow F_{3,\bullet} \rightarrow 0.$$

Denote by $H_{i,\bullet}$ the complex obtained by applying $\text{Hom}(-, N)$ to $F_{i,\bullet}$, so we have

$$H_{i,j} = \text{Hom}(F_{i,-j}, N).$$

For every i the exact sequence

$$0 \rightarrow F_{1,i} \rightarrow F_{2,i} \rightarrow F_{3,i} \rightarrow 0$$

is split exact because $F_{3,i}$ is a free R -module (see Exercise 2.17), and hence also the induced sequence

$$0 \longrightarrow H_{3,-i} \longrightarrow H_{2,-i} \longrightarrow H_{1,-i} \longrightarrow 0$$

is exact (see Exercise 2.19). Applying Theorem 9.5 to the short exact sequence of complexes of abelian groups

$$0 \longrightarrow H_{3,\bullet} \longrightarrow H_{2,\bullet} \longrightarrow H_{1,\bullet} \longrightarrow 0$$

we obtain a long exact sequence

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_{i+1}(H_{2,\bullet}) & \longrightarrow & H_{i+1}(H_{1,\bullet}) & \longrightarrow & \cdots \\ & & \searrow & & \searrow & & \\ & \longrightarrow & H_i(H_{3,\bullet}) & \longrightarrow & H_i(H_{2,\bullet}) & \longrightarrow & H_i(H_{1,\bullet}) & \longrightarrow & \cdots \\ & & \searrow & & \searrow & & \searrow & & \\ & \longrightarrow & H_{i-1}(H_{3,\bullet}) & \longrightarrow & H_{i-1}(H_{2,\bullet}) & \longrightarrow & \cdots & & \end{array}$$

which, taking into account the vanishing of Ext^i for $i < 0$ and the fact that $\text{Ext}^0 = \text{Hom}$, gives precisely the exact sequence of the theorem. One verifies that this sequence does not depend on the choice of free resolutions. \square

3. Ext^1 and extensions

DEFINITION 11.6. Let R be a ring and let M and N be R -modules. An *extension of M by N* is a short exact sequence

$$0 \longrightarrow N \longrightarrow E \longrightarrow M \longrightarrow 0$$

of R -modules. Two such extensions are called *equivalent* if there exists a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & N & \longrightarrow & E & \longrightarrow & M & \longrightarrow & 0 \\ & & \downarrow \text{id} & & \downarrow & & \downarrow \text{id} & & \\ 0 & \longrightarrow & N & \longrightarrow & E' & \longrightarrow & M & \longrightarrow & 0 \end{array}$$

of R -modules. We define the set $\text{ext}_R(M, N)$ to be the set of equivalence classes of extensions of M by N .

Note that the morphism $E \rightarrow E'$ in the above diagram is automatically an isomorphism, see Exercise 2.5. Warning: there can be non-equivalent extensions with $E \cong E'$, see Exercise 11.9.

We will now construct a map

$$\theta: \text{ext}_R(M, N) \rightarrow \text{Ext}_R^1(M, N)$$

and show that it is a bijection. To define the map, consider an element $e \in \text{ext}_R(M, N)$, represented by a short exact sequence

$$0 \longrightarrow N \longrightarrow E \longrightarrow M \longrightarrow 0.$$

By Theorem 11.5 this induces a long exact sequence, of which a part reads

$$\cdots \longrightarrow \text{Hom}_R(N, N) \longrightarrow \text{Ext}_R^1(M, N) \longrightarrow \cdots$$

We define $\theta(e) \in \text{Ext}_R^1(M, N)$ to be the image of id_N under the above map.

THEOREM 11.7. *The map*

$$\theta: \text{ext}_R(M, N) \rightarrow \text{Ext}_R^1(M, N)$$

is a bijection.

PROOF. We construct map

$$\psi: \text{Ext}_R^1(M, N) \rightarrow \text{ext}_R(M, N)$$

but omit the tedious verification that it is a two-sided inverse.

Choose a free module F and a surjection $F \rightarrow M$. Let K be the kernel. Then we have a short exact sequence

$$0 \longrightarrow K \xrightarrow{\gamma} F \longrightarrow M \longrightarrow 0.$$

The induced long exact sequence of Theorem 11.5 contains

$$\text{Hom}_R(F, N) \longrightarrow \text{Hom}_R(K, N) \xrightarrow{\delta} \text{Ext}_R^1(M, N) \longrightarrow \text{Ext}_R^1(F, N).$$

Since F is a free module, it has a ‘trivial’ one-term free resolution, and one sees that $\text{Ext}_R^i(F, N) = 0$ for all $i > 0$. It follows that the map δ is surjective.

Let $e \in \text{Ext}_R^1(M, N)$. Choose an R -linear map $f: K \rightarrow N$ with $\delta(f) = e$. In the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \xrightarrow{\text{id}} & K & \longrightarrow & 0 \\ & & \downarrow & & \downarrow (f, \gamma) & & \downarrow \gamma \\ 0 & \longrightarrow & N & \xrightarrow{\iota_1} & N \oplus F & \xrightarrow{\pi_2} & F \longrightarrow 0 \end{array}$$

both rows are exact, and all the vertical maps are injective (since γ is injective). It follows that there is an induced short exact sequence of cokernels

$$0 \longrightarrow N \longrightarrow E \longrightarrow M \longrightarrow 0$$

with $E = \text{coker}(K \rightarrow N \oplus F)$. We define $\psi(e) \in \text{ext}_R(M, N)$ to be this extension. One verifies that this is well-defined: if also $f': K \rightarrow N$ satisfies $\delta(f') = e$, leading to an extension E' , then $f' = f + h$ for some linear map $h: F \rightarrow N$, and one shows that the isomorphism

$$F \oplus N \rightarrow F \oplus N, (x, y) \mapsto (x, y + hx)$$

induces an isomorphism $E \rightarrow E'$ of extensions. □

Exercises

EXERCISE 11.1. Let K be a field. Let M and N be K -modules. Show that $\text{Ext}_K^i(M, N) = 0$ for all $i \neq 0$.

EXERCISE 11.2. Let n and m be positive integers. Compute for all i the \mathbf{Z} -modules

- (1) $\text{Ext}_{\mathbf{Z}}^i(\mathbf{Z}, \mathbf{Z}/m\mathbf{Z})$;
- (2) $\text{Ext}_{\mathbf{Z}}^i(\mathbf{Z}/n\mathbf{Z}, \mathbf{Z})$;
- (3) $\text{Ext}_{\mathbf{Z}}^i(\mathbf{Z}/n\mathbf{Z}, \mathbf{Z}/m\mathbf{Z})$.

EXERCISE 11.3. Let $R = K[X^2, X^3] \subset K[X]$. Let $I = (X^2, X^3)$. Show that $\text{Ext}_R^1(I, R/I)$ is non-zero, and conclude that I is not a principal ideal. Show that also $\text{Ext}_R^2(R/I, R/I)$ is non-zero.

EXERCISE 11.4. Let R be a principal ideal domain. Show that for every $i \geq 2$, for every finitely generated R -module M , and for every R -module N we have $\text{Ext}_R^i(M, N) = 0$.

EXERCISE 11.5. Let R be a ring, N an R -module and $n \geq 1$. Assume that $\text{Ext}_R^n(M, N) = 0$ for all R -modules M . Show that also $\text{Ext}_R^{n+1}(M, N) = 0$ for all R -modules M . (Hint: consider a short exact sequence of the form $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ with F a free module).

EXERCISE 11.6. Let

$$0 \longrightarrow N \longrightarrow E \longrightarrow M \longrightarrow 0$$

be a short exact sequence of R -modules. Let $\varphi: N \rightarrow N'$ be a morphism of R -modules, and let E' be the push-out of $N \rightarrow E$ and $\varphi: N \rightarrow N'$. Show that there is a short exact sequence

$$0 \longrightarrow N' \longrightarrow E' \longrightarrow M \longrightarrow 0$$

of R -modules.

EXERCISE 11.7. Let

$$0 \longrightarrow N \longrightarrow E \longrightarrow M \longrightarrow 0$$

be a short exact sequence of R -modules. Let $\varphi: M' \rightarrow M$ be a morphism of R -modules, and let E' be the pull-back of $E \rightarrow M$ and $\varphi: M' \rightarrow M$. Show that there is a short exact sequence

$$0 \longrightarrow N \longrightarrow E' \longrightarrow M' \longrightarrow 0$$

of R -modules.

EXERCISE 11.8. Let R be a ring. Show directly (without using the relation with Ext^1) that

$${}_R\mathbf{Mod}^{\text{opp}} \times {}_R\mathbf{Mod} \rightarrow \mathbf{Set}, (M, N) \mapsto \text{ext}_R(M, N)$$

is a functor. (In particular: explain what the functor does on the level of morphisms).

EXERCISE 11.9. Describe explicitly the n elements of the set $\text{ext}_{\mathbf{Z}}(\mathbf{Z}/n\mathbf{Z}, \mathbf{Z})$ for:

- (1) n a prime number;
- (2) $n = pq$ with p and q distinct primes;
- (3) $n = 4$.

EXERCISE 11.10. For $i = 1, 2$ let

$$0 \longrightarrow N \xrightarrow{\alpha_i} E_i \xrightarrow{\beta_i} M \longrightarrow 0$$

be a short exact sequence of R -modules. Define an R -module E as the quotient

$$E := \frac{\ker(\beta_1 - \beta_2: E_1 \oplus E_2 \rightarrow M)}{\text{im}((\alpha_1, \alpha_2): N \rightarrow E_1 \oplus E_2)}.$$

Show that there is a natural short exact sequence

$$0 \longrightarrow N \longrightarrow E \longrightarrow M \longrightarrow 0$$

of R -modules.

EXERCISE 11.11. Let K be a field and let $\lambda_1, \lambda_2 \in K$. Let $M_i := K[X]/(X - \lambda_i)$. Show that $\text{Ext}_{K[X]}^1(M_2, M_1) = 0$ if $\lambda_1 \neq \lambda_2$. Compute $\text{Ext}_{K[X]}^1(M_2, M_1)$ when $\lambda_2 = \lambda_1$.

EXERCISE 11.12. Let K be a field, $\lambda_1, \lambda_2, \mu \in K$. Consider the matrix

$$A := \begin{pmatrix} \lambda_1 & \mu \\ 0 & \lambda_2 \end{pmatrix}$$

and let E be the $K[X]$ -module given by $E = K^2$ on which X acts by A . Show that E sits in a short exact sequence of $K[X]$ -modules

$$0 \longrightarrow M_1 \longrightarrow E \longrightarrow M_2 \longrightarrow 0$$

where $M_i = K$ with X acting as λ_i . Show that this sequence splits if and only if $\lambda_2 \neq \lambda_1$ or $\mu = 0$. Relate this to the previous exercise.

EXERCISE 11.13. Let n be a positive integer, and consider the ring $R := \mathbf{Z}[X]/(X^n - 1)$. Let M be the quotient module $R/(X - 1)$. Compute the R -modules $\text{Ext}_R^i(M, M)$ and $\text{Ext}_R^i(M, R)$. (Hint: use the free resolution from Exercise 10.4).

Bibliography

- [1] ATIYAH, M.F. & MACDONALD, I.G. *Introduction To Commutative Algebra*, Westview Press, 1969.
- [2] LANG, S. — *Algebra*, Graduate Texts in Mathematics 211, Springer, 2002.
- [3] MOERDIJK, I. — *Notes on Homological Algebra*, Lecture notes, 2008.
- [4] STEVENHAGEN, P. — *Algebra 2*, <http://websites.math.leidenuniv.nl/algebra/algebra2.pdf>.
- [5] LEINSTER, T. — *Basic Category Theory*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2014.